

COMMENT

Open Access



Association of cord blood asprosin concentration with atherogenic lipid profile and anthropometric indices

Hanan Khudhair Hussein¹, Malik Aubead², Hamzah H. Kzar³, Yasir Salam Karim⁴, Ali H. Amin^{5,6}, Moaed E. Al-Gazally⁷, Tousief Irshad Ahmed^{8*}, Mohammed Abed Jawad⁹, Ali Thaeer Hammid¹⁰, Abduladheem Turki Jalil^{11,12}, Yasser Fakri Mustafa¹³, Marwan Mahmood Saleh¹⁴ and Hafez Heydari^{15*}

Abstract

Background: Elevated lipids in umbilical cord blood affect fetal programming, leading to a higher risk of developing cardiovascular disease in later life. However, the causes of changes in the lipid profile of umbilical cord blood are not clear yet. This study aimed for the first time to determine the association of asprosin concentration with TAG, TC, HDL-C, LDL-C concentrations and TAG/HDL-C, TC/HDL-C, LDL-C/HDL-C and non-HDL-C/HDL-C ratio in umbilical cord blood as well as newborn anthropometric indices. This cross-sectional study was based on 450 mother- newborn pairs of a birth cohort study in Sabzevar, Iran. Multiple linear regression was used to estimate the association of lipid concentration and lipid ratios as well as birth weight (BW), birth length (BL), head circumference (HC) and chest circumference (CC) with asprosin in cord blood samples controlled for the relevant covariates.

Result: In fully adjusted models, each 1 ng/mL increase in asprosin was associated with 0.19 (95% CI 0.06, 0.31, $P < 0.01$), 0.19 (95% CI 0.10, 0.29, $P < 0.01$), 0.17 (95% CI 0.09, 0.25, $P < 0.01$), 0.17 (95% CI 0.09, 0.25, $P < 0.01$), 0.01 (95% CI 0.00, 0.013, $P < 0.01$), 0.01 (95% CI 0.01, 0.01, $P < 0.01$), 0.01 (95% CI 0.01, 0.01, $P < 0.01$) and 0.01 (95% CI 0.01, 0.01, $P < 0.01$) increase in TAG, TC, LDL-C, TAG/HDL-C, TC/HDL-C, LDL-C/HDL-C and non-HDL-C/HDL-C ratio respectively. Moreover, higher asprosin levels was positively associated with newborn BW, BL, HC and CC; however, these associations were not statistically significant.

Conclusion: Overall, our findings support the positive association between cord asprosin concentration and the development of atherogenic lipid profile in newborns. Further studies are needed to confirm the findings of this study in other populations.

Keywords: Asprosin, Lipid profile, Umbilical cord blood, Triglyceride, Cholesterol

Introduction

Coronary vascular disease (CVDs) is the leading cause of mortality in the world, accounting by 31.5% of all global death [1]. Atherosclerosis is the most common

cause of CVDs, including ischemic heart disease, heart failure, and peripheral arterial disease [2, 3]. The atherogenic processes involved in CVDs pathogenesis begin in the intrauterine environment during fetal development and then gradually progress in subsequent years [4]. By detecting CVDs lipid risk factors including Apo A1, Apo B and atherogenic index in the umbilical cord blood of the term newborns, it is possible to recognize newborns at a higher risk for CVDs in the future [5].

*Correspondence: khagankhan@gmail.com; hafezheydari62@yahoo.com

⁸ Department of Biochemistry, SKIMS, Srinagar, J&K, India

¹⁵ Non-Communicable Diseases Research Center, Sabzevar University of Medical Sciences, Sabzevar, Iran

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

The causes of changes in lipid metabolism of umbilical cord blood are not completely known. Still, there is some evidence that maternal adiposity status and adipokines (including resistin and leptin) may play an active role in regulating fetal lipid profile [6–13]. Asprosin, a novel adipokine identified by Omre et al., is secreted in response to starvation from adipocytes and promotes rapid releases of glucose from the liver cells [14]. The association between circulating asprosin and lipids metabolism has been identified [15, 16]. Asprosin was positively correlated with LDL-c, APOB, APOE and TAG concentrations and negatively correlated with HDL-C concentration [15–18]. Therefore, it can be speculated that asprosin may be involved in a process that develops an atherogenic lipid profile in umbilical cord blood. To the best of our knowledge, there is no study on the association of asprosin and umbilical cord blood lipid profile. Therefore, this study aimed to investigate the relationship between umbilical cord blood asprosin with lipid profile and newborn anthropometric indices. Besides, the relationship between umbilical asprosin with different lipid ratios as predictors of CVDs has been studied.

Material and methods

Study population

This study was conducted from June 2018 to August 2018 at the Mobini Hospital in Sabzevar (a town in the Khorasan-Razavi province), Iran. We enrolled 450 healthy pregnant women recruited to the hospital for their delivery based on our inclusion/exclusion criteria. The inclusion criteria were mothers with a normal term of pregnancy (37–42 weeks of gestation), normal singleton pregnancies, normal vaginal delivery, normal lipid profile, self-report of no hypertension, preeclampsia, diabetes, and gestational diabetes. The exclusion criteria included intrauterine fetal growth restriction, fetal structural abnormalities, and drug intake that affects metabolism. It should be noted that the inclusion/exclusion criteria were detected through self-reports and medical records of participants. The project was approved by the Clinical Research Ethical Committee (IR.MEDSAB.REC.1397.012) of Sabzevar University of Medical Sciences and all participants signed the informed consent form before to enrollment.

Cord blood collection and biochemical assessments

Cord blood samples were collected from the umbilical cord vein in serum separator tubes with a clot activator at delivery time. Sera were separated, aliquoted and stored at -80 until subsequent analysis. TAG, HDL-C, and TC were measured using an automated analyzer (Biotechnica, BT 1500, Rome, Italy) and commercial kits (P.L:38,231,049, Pars Azmoon, Tehran, Iran) by

enzymatic colorimetric methods. LDL-C concentration was calculated using the Friedewald equation. Asprosin concentration was quantified by enzyme-linked immunosorbent assay (ELISA) using a commercially available kit according to manufacture instructions (Cat. No: CK-E91570, EASTBIOPHARM, China). Asprosin concentration was reported in ng/mL. The limit of detection (LOD) was 0.34 ng/mL. The intra-assay and interassay coefficients of variation were < 10% and < 12%, respectively.

Statistical analyses

Main analysis

All statistical analyses were performed using Stata version 15 software (Stata Corp LP, College Station, Texas). The Shapiro–Wilk test and P-P plot tested the distribution of data. Data normally distributed were shown as mean \pm SD and data with skewed distribution were shown as median (interquartile range (IQR)).

Multiple regression models were used to investigate the association of umbilical asprosin concentration and lipid profile and anthropometric indices. Models were further adjusted for a *priori* of the selected variables: maternal age and pre-pregnancy BMI, parity, gestational age at delivery, environmental tobacco exposure at home during pregnancy (Yes/No) and two household socioeconomic status (SES) indicators as well as two neighborhood SES indicators. Maternal education (no education, primary school, secondary school or university) and income (15 million, 15–30 million and more than 30 million riyals) were used as household SES indicators, and illiterate adults and unemployment percentages per census tract based on the last census of Iran (2016) were used as neighborhood SES indicators. Regression coefficients were reported for each 1 ng/ml increase in asprosin concentration. A significant level of 0.05 was used for all statistical analyses.

Sensitivity analysis

Asprosin is introduced as a fasting-induced hormone and the level of asprosin in cord blood may be significantly determined by the glucose concentration [19]. Therefore we further adjusted our models for umbilical blood glucose. Moreover, we further adjusted our models with the mother's lipid profile during pregnancy.

Results

Descriptive statistics

The descriptive statistics of the participants are presented in Table 1. The mean (standard deviation (SD)) age of mothers was 27.65 (5.36) years. Of the 450 newborns included in our study, 225 (50%) were male. The median (IQR) of umbilical TAG, TC, LDL-C, HDL-C, were 29.9

Table 1 Descriptive statistics of participants and lipid profile

Parameter	Discription
Maternal characteristics	
Age (year); mean (SD)	27.7 (5.4)
Pre-pregnancy BMI (kg/m ²); Median (IQR)	24.6 (5.69)
Parity; Median (IQR)	2 (1)
Gestational age (day); Median (IQR)	278 (12)
Birth weight (gr); Median (IQR)	3250 (550)
Birth length (cm); Median (IQR)	51 (3)
Head circumference (cm); Median (IQR)	35 (2)
Chest circumference (cm); Median (IQR)	33 (3)
Education	
No education/ primary school; N (%)	169 (37.6)
Secondary school education; N (%)	212 (47.1)
University degree or higher; N (%)	69 (15.3)
Self-reported tobacco exposure at home	
Yes; N (%)	81 (18.0)
No; N (%)	369 (82.0)
Income	
≥ 15 million Rials; N (%)	221 (49.1)
15 to 30 million Rials; N (%)	100 (22.2)
30 ≤ million Rials; N (%)	79 (17.5)
Illiterate percent per census tract (%); Median (IQR)	25.38 (17.5)
Unemployed percent per census tract (%); Median (IQR)	6.91 (5.1)
TAG (mg/dl); Median (IQR)	29.9 (26.9)
TC (mg/dl); Median (IQR)	60.4 (28)
HDL-C (mg/dl); Median (IQR)	28.2 (12.5)
LDL-C (mg/dl); Median (IQR)	23.28 (21.48)
Non-HDL-C (mg/dl); Median (IQR)	30.8 (21)
TC/HDL-C; Median (IQR)	2.01 (0.76)
TAG/HDL-C; Median (IQR)	1.15 (1.18)
LDL-C/HDL-C; Median (IQR)	0.78 (0.67)
Non-HDL-C/HDL-C; Median (IQR)	1.01 (0.76)
Asprosin (ng/mL); Median (IQR)	30.44 (19.08)

(26.9), 60.4 [27], 23.28 (21.48), 28.2 (12.5) mg/dL, respectively. Moreover, the median (IQR) for TAG/ HDL-C, TC /HDL-C, and Non-HDL-C/HDL-C were 1.15 (1.18), 2.01 (0.766) and 1.01 (0.76), respectively. The median (IQR) of umbilical asprosin was 30.44 (19.08) ng/mL. No significant differences were observed in asprosin concentration as well as lipid profile between male and female newborns (P -value > 0.05). The median (IQR) for birth weight (BW), birth length (BL), head circumference (HC) and chest circumference (CC) were 3250 (550) gr, 51 (3) cm, 35 (2) cm and 33 (3) cm, respectively.

Association with anthropometric indices

The results of the relationship between umbilical cord blood concentration of asprosin and BW, BL, HC and CC

Table 2 Association of cord blood asprosin concentration and anthropometric indices

Anthropometric indices	Model	β-coefficient (95% CI)	P value
Birth weight	Crude	0.64 (−1.21, 2.50)	0.49
	Adjusted	0.39 (−1.55, 2.32)	0.69
Birth length	Crude	0.01 (−0.01, 0.02)	0.42
	Adjusted	0.00 (−0.01, 0.02)	0.59
Head circumference	Crude	0.00 (−0.01, 0.01)	0.74
	Adjusted	0.00 (−0.01, 0.01)	0.52
Chest circumference	Crude	0.02 (−0.03, 0.07)	0.39
	Adjusted	0.01 (−0.06, 0.07)	0.29
Gestational age at delivery*	Crude	0.03 (−0.01, 0.06)	0.13
	Adjusted	0.02 (−0.02, 0.06)	0.24

Adjusted for Age of mother, BMI of mother before pregnancy, Number of pregnancies, Gestational age, Percent of illiterate per census tract, Percent of unemployment per census tract, Paternal education, Maternal education, Income and Tobacco exposure at home. The regression coefficient was reported based on 1 ng/mL increase in asprosin concentration

CI confidence interval

* Adjusted for above covariates except for Gestational age

are presented in Table 2. In crude and adjusted models, higher levels of cord blood asprosin were positively associated with BW, BL, HC and CC as well as gestational age at delivery; however, these associations were not statistically significant.

Association with lipid biomarkers

The association of asprosin concentration and lipid profile are presented in Table 3. In fully adjusted models increase in umbilical asprosin concentration was associated with increase in umbilical TAG, TC, LDL-C levels as well as TAG/HDL-C, TC/HDL-C, LDL-C/HDL-C and non-HDL-C/HDL-C ratio. Each 1 ng/ml increase in asprosin was associated with 0.19 mg/dL (95% CI 0.06, 0.31, P <0.01), 0.19 mg/dL (95% CI 0.10, 0.29, P <0.01), 0.17 mg/dL (95% CI 0.09, 0.25, P <0.01), 0.17 (95% CI 0.09, 0.25, P <0.01), 0.01 (95% CI 0.00, 0.01, P <0.01), 0.01 (95% CI 0.01, 0.01, P <0.01), 0.008 (95% CI 0.01, 0.01, P <0.01) and 0.01 (95% CI 0.01, 0.01, P <0.01) increase in TAG, TC, LDL-C, TAG/HDL-C, TC/HDL-C, LDL-C/HDL-C and non-HDL-C/HDL-C ratio respectively. The association of asprosin concentration and HDL-C concentration was not statistically significant (P -value = 0.43).

Sensitivity analysis

The results of adjusted our models for umbilical blood glucose and mother's lipid profile on the association of asprosin concentration with anthropometric indices and umbilical lipid profile showed no notable differences in

Table 3 Linear regression for cord asprosin concentration and change in lipid profile in newborns

		β -coefficient (95% CI)	P_value
Lipid profile			
	TAG	Crude 0.19 (0.08, 0.3)	<0.01
	Adjusted	0.19 (0.07, 0.32)	<0.01
TC	Crude	0.19 (0.11, 0.28)	<0.01
	Adjusted	0.20 (0.10, 0.29)	<0.01
LDL-C	Crude	0.17 (0.10, 0.24)	<0.01
	Adjusted	0.18 (0.10, 0.26)	<0.01
HDL-C	Crude	-0.02 (-0.06, 0.02)	0.36
	Adjusted	-0.02 (-0.06, 0.03)	0.43
Lipid ratio			
	TAG/HDL-C	Crude 0.01 (0.00, 0.01)	<0.01
	Adjusted	0.01 (0.00, 0.01)	<0.01
TC/HDL-C	Crude	0.01 (0.01, 0.01)	<0.01
	Adjusted	0.01 (0.01, 0.01)	<0.01
LDL-C/HDL-C	Crude	0.01 (0.01, 0.01)	<0.01
	Adjusted	0.01 (0.01, 0.01)	<0.01
Non-HDL-C/HDL-C	Crude	0.01 (0.01, 0.01)	<0.01
	Adjusted	0.01 (0.01, 0.01)	<0.01

Adjusted for Age of mother, BMI of mother before pregnancy, Number of pregnancies, Gestational age, Percent of illiterate per census tract, Percent of unemployment per census tract, Paternal education, Maternal education, Income and Tobacco exposure at home. The regression coefficient was reported based on 1 ng/mL increase in asprosin concentration

CI confidence interval

the main results in terms of significance and direction (data not shown).

Discussion

To the best of our knowledge, this is the first study on the relationship between asprosin concentration with some anthropometric indices (e.g., HC and CC) and lipid profile in cord blood samples. We observed that serum asprosin concentration was positively associated with TAG, TC, LDL-C concentrations and TAG/HDL-C, TC/HDL-C, LDL-C/HDL-C and non-HDL-C/HDL-C ratio in umbilical cord blood. However, we did not find any significant association with HDL-C concentrations in umbilical cord blood. Moreover, there was a positive but not statistically significant association between asprosin levels and anthropometric indices.

Interpret the results

We did not find a study on the association of asprosin concentration and lipid profile in cord blood; therefore, it is impossible to compare our results with other studies. However, several studies have examined the association of serum asprosin concentrations and lipid profile in adults. Wang et al. 2018 showed that serum

concentrations of asprosin were positively correlated with TAG concentrations. They also reported that asprosin concentrations were negatively correlated with HDL-C concentrations [20]. Moreover, the association of serum concentration of asprosin and HDL-C has been reported in a study by Long et al. [16]. The studies by Whang et al. 2018 and Zhang et al. 2019 reported there was no statistically significant association between serum concentrations of asprosin and LDL-C [15, 20]. However, Li et al. 2018 showed that serum concentrations of asprosin were positively associated with LDL-C, apoB and apoE [21].

We did not observe any significant difference in lipid profile as well as asprosin concentration between male and female newborns. However, a study by Kelishadi et al. 2007 reported that total and HDL cholesterol in girls were significantly higher than in boys [9]. Moreover, a study by Ghiasi et al. 2013 reported that the cord blood from female newborns had higher levels of LDL-C, HDL-C and total cholesterol compared to male newborns [22]. Furthermore, Jahanfar et al. 2016 in a cohort study of twins, reported that male-male twins were heavier than male-females and female-female twin pairs [23].

Our study found a positive but non-significant association between asprosin concentration and anthropometric indices. A study by Hoffmann et al. 2022 on 247 umbilical plasma samples reported that there was no significant correlation between BW and BL with asprosin concentration [24].

Biological plausibility

Asprosin may affect lipid metabolism by two different mechanisms: the effect on the insulin signaling pathway and proopiomelanocortin (POMC)-positive neurons activity. Some effects of asprosin on lipid profile may be exerted by altering gene expression in liver cells through insulin resistance induction [25–28]. High insulin concentrations in IR alter the expression of some genes [29–32]. In the liver, the target tissue of asprosin, insulin induces the expression and activation of enzymes involved in hepatic lipid production [33–37]. Although the molecular mechanism of asprosin on insulin signaling is not well known; however, several studies suggested elevated concentrations of asprosin interfere with insulin function and induces insulin resistance [38–41]. Romere et al. 2016 showed that single injection of asprosin results in hyperglycemia and hyperinsulinemia [14]. Wang et al. 2018 found a negative relationship between serum asprosin concentrations and indicators regarding the first-phase insulin secretion, such as AUC, AIR, and GDI and HOMA- β , and a positive correlation with HOMA-IR [20]. They suggested that the asprosin-related metabolic pathways dysregulation might be through its role in β -cell dysfunction and insulin resistance [20]. Jung et al. 2019

showed that treatment of skeletal muscle cells by recombinant asprosin results in impairment of insulin sensitivity through PKC δ -associated ER stress/inflammation pathways [26]. Lee et al. 2019 showed that treatment of primary human islets with recombinant asprosin induced the inflammation response, cellular dysfunction, and apoptosis in a dose-dependent manner [42].

Asprosin exerts some of its effects by changing the function of the CNS system. Duerrschmid et al. 2017 demonstrate that asprosin in the circulation crosses the blood–brain barrier and inhibits the downstream anorexigenic proopiomelanocortin (POMC)-positive neurons. The inhibition of these neurons leads to appetite stimulation and a drive to accumulate adiposity and body weight [19, 43, 44]. Blockade of the CNS melanocortin system increased triglyceride synthesis in the periphery and triglyceride content in the liver [45, 46].

Limitations

Our study has some limitations that should be considered in future studies. The sample size was relatively limited, and the findings of this study required to confirm from other ethnicities. Also, the cross-sectional nature of this study did not allow us to address the underlying signaling pathways. Moreover, we did not have any data on pre-pregnancy lipid profile, maternal diet, maternal weight gain and other prevalent endocrine disorders like PCOS that can affect our results. Furthermore, we missed data about timing of cord clamping that could affect TG and total cholesterol. Research in cellular models is required to determine underlying mechanisms.

Conclusion

The present study indicated that asprosin concentration in umbilical cord blood was positively associated with TAG, TC, LDL-C concentration and TAG/HDL-C, TC/HDL-C, LDL-C/HDL-C and non-HDL-C/HDL-C ratio. Our data provide new clinical information about the role of adipokines in the regulation of atherogenic lipids including TC and LDL-C in umbilical cord blood. Further studies including the identification of the molecular changes in tissue structure or functions, are needed for a better understanding of asprosin- lipid profile and its clinical outputs in humans.

Acknowledgements

The authors thank form the staff of Shahidan-Mobini Hospital and Sabzevar University of Medical Sciences for their help and support. Moreover, the authors would like to thank the Deanship of Scientific Research at Umm Al-Qura University for supporting this work by Grant Code: (22UQU4331100DSR05).

Author contributions

HKH: Conceptualization, Methodology, Software, Writing- Original draft preparation. MA: Data curation, Writing—Original draft preparation. HHK: Visualization, Investigation. YSK: Visualization, Investigation., AHA: Visualization,

Investigation., MEA: Visualization, Investigation., TIA: Visualization, Investigation., MAJ: Visualization, Investigation. ATH: Visualization, Investigation Writing-Reviewing and Editing. ATJ: Visualization, Investigation, Writing—Reviewing and Editing. YFM: Supervision, Writing—Reviewing and Editing. MMS: Supervision, Software, Validation, Writing—Reviewing and Editing. HH, Writing—Original Draft Preparation, Writing—Review & Editing. All authors read and approved the manuscript.

Funding

None.

Availability of data and materials

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Declarations

Ethics approval and consent to participate

The project was approved by the Clinical Research Ethical Committee (IR. MEDSAB.REC.1397.012) of Sabzevar University of Medical Sciences and all participants signed the informed consent form prior to enrollment. All authors confirm that all methods were carried out in accordance with relevant guidelines.

Consent for publication

Not applicable.

Competing interests

The authors declared they do not have anything to disclose regarding conflict of interest with respect to this manuscript.

Author details

¹College of Medicine, University of Babylon, Babil, Iraq. ²Department of Obstetrics and Gynecology, Hammurabi College of Medicine, University of Babylon, Babil, Iraq. ³Veterinary Medicine College, Al-Qasim Green University, Al-Qasim, Iraq. ⁴Al-Manara College For Medical Sciences, Maysan, Iraq. ⁵Deanship of Scientific Research, Umm Al-Qura University, Makkah, Saudi Arabia. ⁶Zoology Department, Faculty of Science, Mansoura University, Mansoura, Egypt. ⁷College of Medicine, University of Al-Ameed, Karbala, Iraq. ⁸Department of Biochemistry, SKIMS, Srinagar, J&K, India. ⁹Al-Nisour University College, Baghdad, Iraq. ¹⁰Computer Engineering Techniques, Faculty of Information Technology, Imam Ja'afar Al-Sadiq University, Baghdad, Iraq. ¹¹Faculty of Biology and Ecology, Yanka Kupala State University of Grodno, 230023 Grodno, Belarus. ¹²College of Technical Engineering, The Islamic University, Najaf, Iraq. ¹³Department of Pharmaceutical Chemistry, College of Pharmacy, University of Mosul, Mosul 41001, Iraq. ¹⁴Department of Biophysics, College of Applied Sciences, University Of Anbar, Ramadi, Iraq. ¹⁵Non-Communicable Diseases Research Center, Sabzevar University of Medical Sciences, Sabzevar, Iran.

Received: 26 March 2022 Accepted: 5 May 2022

Published online: 18 May 2022

References

- Benjamin EJ. Correction to: heart disease and stroke statistics-2017 update: a report from the American Heart Association. *Circulation*. 2017;135(10):e646.
- Herrington W, Lacey B, Sherliker P, Armitage J, Lewington S. Epidemiology of atherosclerosis and the potential to reduce the global burden of atherothrombotic disease. *Circ Res*. 2016;118(4):535–46.
- Salari A, Roshanaei K, Rasoulian B, Khalili Fard J. Carvacrol attenuates disrupted lipid profile induced by organophosphates in male wistar rat: a comparative toxicity. *J Chem Health Risks*. 2021;11(Special Issue: Bioactive Compounds: Their Role in the Prevention and Treatment of Diseases):121–8.
- Napoli C, D'Armiento FP, Mancini FP, Postiglione A, Witztum JL, Palumbo G, et al. Fatty streak formation occurs in human fetal aortas and is greatly enhanced by maternal hypercholesterolemia. Intimal accumulation of

- low density lipoprotein and its oxidation precede monocyte recruitment into early atherosclerotic lesions. *J Clin Invest.* 1997;100(11):2680–90.
5. Casanueva V, Cid X, Chiang M, Molina M, Ferrada M, Perez R, et al. Lipids, lipoproteins and apolipoproteins in normal newborns. *Rev Med Chil.* 1998;126(9):1073–8.
 6. Nelson SM, Freeman DJ, Sattar N, Johnstone FD, Lindsay RS. IGF-1 and leptin associate with fetal HDL cholesterol at birth: examination in offspring of mothers with type 1 diabetes. *Diabetes.* 2007;56(11):2705–9.
 7. Fonseca MJ, Santos AC. Umbilical cord blood adipokines and newborn weight change. *Arch Gynecol Obstet.* 2015;291(5):1037–40.
 8. Xiao-yun C. A study of changes of leptin, cholesterol and triglyceride in cord blood of newborns. *J Taishan Med Coll.* 2008;10.
 9. Kelishadi R, Badiiee Z, Adeli K. Cord blood lipid profile and associated factors: baseline data of a birth cohort study. *Paediatr Perinat Epidemiol.* 2007;21(6):518–24.
 10. Khan W, Augustine D, Rao RS, Patil S, Awan KH, Sowmya SV, et al. Lipid metabolism in cancer: a systematic review. *J Carcinog.* 2021;20:4.
 11. Sowndarya K, Joseph JA, Shenoy A, Hegde A. Evaluation of triglyceride/high-density lipoprotein ratio as a surrogate marker for insulin resistance in healthy young males. *J Nat Sci Biol Med.* 2021;12:213.
 12. Vaccaro C, Shakeri A, Czaplinski E, Eltonsy S. New-generation antiepileptic drugs during pregnancy and the risk of attention-deficit hyperactivity disorder: a scoping review. *J Popul Ther Clin Pharmacol.* 2020;27(4):e1–18.
 13. Nakhaei P, Margiana R, Bokov DO, Abdelbasset WK, Kouhbanani MAJ, Varma RS, et al. Liposomes: structure, biomedical applications, and stability parameters with emphasis on cholesterol. *Front Bioeng Biotechnol.* 2021;9:705886.
 14. Romere C, Duerrschmid C, Bournat J, Constable P, Jain M, Xia F, et al. Asprosin, a fasting-induced glucogenic protein hormone. *Cell.* 2016;165(3):566–79.
 15. Zhang L, Chen C, Zhou N, Fu Y, Cheng X. Circulating asprosin concentrations are increased in type 2 diabetes mellitus and independently associated with fasting glucose and triglyceride. *Clin Chim Acta.* 2019;489:183–8.
 16. Long W, Xie X, Du C, Zhao Y, Zhang C, Zhan D, et al. Decreased circulating levels of asprosin in obese children. *Horm Res Paediatr.* 2019;91(4):271–7.
 17. Rerksuppaphol L, Rerksuppaphol S. Comparison of equations for the calculation of low-density lipoprotein cholesterol in Thai population. *J Nat Sci Biol Med.* 2021;12:224.
 18. j Manna M, Jabur MS. The value of lisinopril in managing impaired fertility in experimental rats following alloxan induced hyperglycemia. *J Pharm Negat Results.* 2022;12(2):18.
 19. Duerrschmid C, He Y, Wang C, Li C, Bournat JC, Romere C, et al. Asprosin is a centrally acting orexigenic hormone. *Nat Med.* 2017;23(12):1444–53.
 20. Wang Y, Qu H, Xiong X, Qiu Y, Liao Y, Chen Y, et al. Plasma asprosin concentrations are increased in individuals with glucose dysregulation and correlated with insulin resistance and first-phase insulin secretion. *Mediators Inflamm.* 2018;2018:9471583.
 21. Li X, Liao M, Shen R, Zhang L, Hu H, Wu J, et al. Plasma asprosin levels are associated with glucose metabolism, lipid, and sex hormone profiles in females with metabolic-related diseases. *Mediators Inflamm.* 2018;2018:7375294.
 22. Ghiasi A, Ziaee S, Faghihzadeh S. A Comparison of lipid and lipoprotein B-100 (Apo B-100) Levels in the umbilical cord blood of male and female newborns and the assessment of their impact on neonatal anthropometric indices. *Pathobiol Res.* 2013;16(1):55–62.
 23. Jahanfar S, Lim K. The impact of gender on anthropometric measures of twins. *Twin Res Hum Genet.* 2016;19(6):652–8.
 24. Hoffmann T, Morcos YAT, Janoschek R, Turnwald E-M, Gerken A, Müller A, et al. Correlation of metabolic characteristics with maternal, fetal and placental asprosin in human pregnancy. *Endocr Connect.* 2022;11(3):e220069.
 25. Alan M, Gurlek B, Yilmaz A, Aksit M, Aslanipour B, Gulhan I, et al. Asprosin: a novel peptide hormone related to insulin resistance in women with polycystic ovary syndrome. *Gynecol Endocrinol.* 2019;35(3):220–3.
 26. Jung TW, Kim HC, Kim HU, Park T, Park J, Kim U, et al. Asprosin attenuates insulin signaling pathway through PKCdelta-activated ER stress and inflammation in skeletal muscle. *J Cell Physiol.* 2019;234(11):20888–99.
 27. Pawitan JA, Leviana M, Sukmawati D, Liem IK, Margiana R, Tarcisia T. Prospect of umbilical cord mesenchymal stem cell culture waste in regenerative medicine. *J Global Pharma Technol.* 2017;9(7):1–5.
 28. Margiana R, Jusuf AA, Lestari SW. Immunohistochemistry Detection Method of Rejection Reaction of Human Umbilical Cord Derived Mesenchymal Stem Cell on Rat Sciatic Nerve Tissue. 2009.
 29. Lebovitz HE. Insulin resistance: definition and consequences. *Exp Clin Endocrinol Diabetes.* 2001;109(Suppl 2):S135–48.
 30. Kalvisa A, Siersbaek MS, Praestholm SM, Christensen LJJ, Nielsen R, Stohr O, et al. Insulin signaling and reduced glucocorticoid receptor activity attenuate postprandial gene expression in liver. *PLoS Biol.* 2018;16(12):e2006249.
 31. Liu Y, Nakagawa Y, Wang Y, Liu L, Du H, Wang W, et al. Reduction of hepatic glucocorticoid receptor and hexose-6-phosphate dehydrogenase expression ameliorates diet-induced obesity and insulin resistance in mice. *J Mol Endocrinol.* 2008;41(2):53–64.
 32. Ahmad K. Insulin sources and types: a review of insulin in terms of its mode on diabetes mellitus. *J Tradit Chin Med.* 2014;34(2):234–7.
 33. Foufelle F, Ferre P. New perspectives in the regulation of hepatic glycolytic and lipogenic genes by insulin and glucose: a role for the transcription factor sterol regulatory element binding protein-1c. *Biochem J.* 2002;366(Pt 2):377–91.
 34. Titchenell PM, Lazar MA, Birnbaum MJ. Unraveling the regulation of hepatic metabolism by insulin. *Trends Endocrinol Metab.* 2017;28(7):497–505.
 35. Santoleri D, Titchenell PM. Resolving the paradox of hepatic insulin resistance. *Cell Mol Gastroenterol Hepatol.* 2019;7(2):447–56.
 36. Cook JR, Langlet F, Kido Y, Accili D. Pathogenesis of selective insulin resistance in isolated hepatocytes. *J Biol Chem.* 2015;290(22):13972–80.
 37. Bahrebar M, Adalatpanah Y. The effect of consumption of the alcoholic extract of cedar on blood glucose, urea and total cholesterol diabetic rats. *J Chem Health Risks.* 2021;11((Special Issue: Bioactive Compounds: Their Role in the Prevention and Treatment of Diseases)):93–7.
 38. Wang M, Yin C, Wang L, Liu Y, Li H, Li M, et al. Serum asprosin concentrations are increased and associated with insulin resistance in children with obesity. *Ann Nutr Metab.* 2019;75(4):205–12.
 39. Groener JB, Valkanou A, Kender Z, Pfeiffenberger J, Kihm L, Fleming T, et al. Asprosin response in hypoglycemia is not related to hypoglycemia unawareness but rather to insulin resistance in type 1 diabetes. *PLoS ONE.* 2019;14(9):e0222771.
 40. Zhang X, Jiang H, Ma X, Wu H. Increased serum level and impaired response to glucose fluctuation of asprosin is associated with type 2 diabetes mellitus. *J Diabetes Investig.* 2019;11(12):349–55.
 41. Pradhan MR, Patel SK, Patel S. Risk Factors and spatial clustering of primary infertility in India. *J Infertil Reprod Biol.* 2021;9(1):27–34.
 42. Lee T, Yun S, Jeong JH, Jung TW. Asprosin impairs insulin secretion in response to glucose and viability through TLR4/JNK-mediated inflammation. *Mol Cell Endocrinol.* 2019;486:96–104.
 43. Millington GW. The role of proopiomelanocortin (POMC) neurones in feeding behaviour. *Nutr Metab.* 2007;4:18.
 44. Mori Y. Regulation of appetite by melanocortin and its receptors. *Nihon Rinsho Japanese J Clin Med.* 2001;59(3):431–6.
 45. Nogueiras R, Wiedmer P, Perez-Tilve D, Veyrat-Durebex C, Keogh JM, Sutton GM, et al. The central melanocortin system directly controls peripheral lipid metabolism. *J Clin Invest.* 2007;117(11):3475–88.
 46. Albarado DC, McClaine J, Stephens JM, Mynatt RL, Ye J, Bannon AW, et al. Impaired coordination of nutrient intake and substrate oxidation in melanocortin-4 receptor knockout mice. *Endocrinology.* 2004;145(1):243–52.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.