

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/344876299>

Effect of Cutting Type and Humic Acid on Growth and Root Traits of Fig (L.) Wazeri cv. Saplings Ficus carica

Article in Indian Journal of Ecology · October 2020

CITATIONS

0

READS

212

5 authors, including:



Ahmed Fatkhan

University of Anbar, Ramadi, Iraq

43 PUBLICATIONS 69 CITATIONS

SEE PROFILE



Ali F Almehemdi

University of Anbar

92 PUBLICATIONS 106 CITATIONS

SEE PROFILE



Mukhalad Hadi Asmail

University of Anbar

26 PUBLICATIONS 6 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



university of anbar at glance [View project](#)



RG progressing researcher Certificate. [View project](#)



Effect of Cutting Type and Humic Acid on Growth and Root Traits of Fig (*Ficus carica* L.) Wazeri cv. Saplings

Mukhalad H.I. Alani, Mohammed Khazaal Hameed, Ahmed F.Z. Al-Dulaimy and Ali F. Almehemdi¹

Department of Horticulture, College of Agriculture, University of Anbar, Anbar, Iraq

¹Center of Desert Studies, University of Anbar, Anbar, Iraq

Email: alimsc1973@gmail.com

Abstract: Greenhouse trial was carried out at University of Anbar, Iraq, during the growing season of 2018 to study the effect of the cutting type (terminal, middle and basal) and the application of humic acid 0, 2 and 4 ml/litter on some traits of vegetative and root growth of fig. The humic acid was added to the cultural media in three stations. The different types of cutting showed a significant effect on all vegetative and root growth traits (except the dry weight of the vegetative system). Basal cuttings resulted in maximum s saplings height (49.86 cm), branches per sapling (7.69), stem diameter (12.82 mm), sapling leaves area (180.5 cm) and root length (41.61 cm), respectively. The addition of 4 ml/ litter concentration gave the maximum saplings height, number of branches, -1stem diameter, sapling leaves area, dry weight of vegetative system, roots length and dry weight of the root system (53.42 cm, 7.58 branch sapling, 13.00 mm, 176.2 cm, 21.30 g, 42.14 cm and 8.43 g, respectively). The basal cuttings resulted in positive effect on vegetative and root traits as compared to the middle and terminal cuttings.

Keywords: Figs, Cutting type, Humic acid, Vegetative growth, Root growth

Fig (*Ficus carica* L.) belongs to the Moraceae family which includes more than 800 species (Harrison 2005). The origin of fig is West Asia, and its cultivation spread in the Mediterranean region (Herre et al 2008). Recently, fig cultivation has spread in different parts of the world such as Turkey, Egypt, Spain, Greece, America, Italy, Brazil (Mars et al 2008). In Iraq, the climate is suitable for figs culturing, but cultivation is not at the commercial level. The figs productivity is estimated as 3104 tonnes in 2018 with decrease of 7.32% from last year production (CSSIT 2018). Fig is propagated by many methods such as the propagation by stem cutting taken from branches of one to many years during the bud dormancy period which is the most common vegetative propagation method (Hartmann et al 2002). There are many factors affecting the propagation by stem cuttings, the important one of these factors is the time of taking and planting of the cutting. Furthermore, choosing the type of cutting is the other most important factor, which is affecting the rooting ability of the cuttings relied on their physiological and anatomical status. Hamooh (2004) and Reddy et al (2008) observed that the hardwood fig cuttings were significantly superior in all root traits compared with semi hardwood cuttings. Khapare et al (2012) showed that hardwood fig cuttings were superior in rooting percentage and all the other studied traits.

Agriculture/Anbar University/Iraq, during the growing season of 2018. Cuttings have been taken on January 20th, 2018 from eight years-old fig. Factorial experiment with two factors (cuttings type and humic acid concentrations) was done in Completely Randomized Block Design (RCBD) with three replicates and 15 cuttings for each experimental unit. Three types of cutting (terminal, middle and basal) were used in this study. These cutting with 15–20 cm lengths were collected from one-year-old branches. The cuttings were treated with the fungicide Benomyl 4 g l⁻¹ for 30 minimums, and then they were fully dried, and then bases dipped for 5 s in 2000 mg l⁻¹ of the auxin IBA. Cuttings were then planted in polyethylene bags of 5 kg capacity that contain soil mixture (3 river sand: 1 peatmoss) (Table 1). This culture medium had been sterilized in 4% formaldehyde on 1st of April. The rooted saplings were translocated to the lath house to avoid damages by high temperature in the greenhouse. Humic acid was added to the bags in three concentrations viz. 0, 2 and 4 ml l⁻¹ on 20th of April, 20th of July and 20th of October. Shoot and root traits were recorded at the end of the experiment on 20th of November, which included, saplings height (cm), number of branches (branch saplings⁻¹), stem diameter (mm), sapling leaves area (cm²), dry weight of vegetative system (g), root length (cm) and dry weight of the root system (g).

MATERIAL AND METHODS

Greenhouse experiment was conducted at College of

RESULTS AND DISCUSSION

Sapling height (cm): The different cutting types significantly

affected in sapling height, basal and middle cuttings (C_2 and C_3) significantly superior with the terminal ones with mean of 49.86 and 49.04 cm). The increase in rooting supporting materials such as auxins and other rooting factors in the basal cuttings could be a reason to be led this result (Owais 2010, Rana and Sood 2012), where the auxin function providing saccharides in lateral root sites due to the increased starch motion by the increased vitality of carbohydrate metabolic enzymes to free energy to form enzymes and enhance cell dividing and disclosing enzymes (Bhattacharya and Nanda 1978). All of what mentioned above have their own participation on the enhancement of transplants shoot and roots growth. Also budding in woody cuttings has gained more maturity and germination and growth ready physiological status. The addition of humic acid @ 4 ml l⁻¹ h gave the highest mean (53.42 cm) compared with the interaction between the two factors showed significant differences (application of 4 ml l⁻¹ humic acid to the middle cuttings (C_2H_4) gave the maximum height (56.44 cm) (Table 2). The positive effect of humic acid addition in enhancement of vegetative and root traits of the saplings, may be because of its role for improving soil chemical, physical and biological properties (Schnitzer and Khan 1971). Furthermore, it contains many plants essential nutrients. On other hands, Atee and Al-Sahae (2007) referred that the humic acid has a role in increasing the cation exchange capacity, reducing soil pH and make macro and micro nutrients more available, as well as building a highly efficient root system to absorb

nutrients, which positively reflected on plant growth (Eissa et al 2007).

Branches number (branches sapling⁻¹): Basal cutting (C_3) gave the highest number of branches per sapling 7.69 branch sapling⁻¹, but not significantly different from the treatment C_2 but was significantly better than with the terminal cuttings (C_1). The increase in rooting supporting materials such as auxins and other rooting factors in the basal cuttings could be a reason to be led this result (Owais 2010). The addition of humic acid resulted in significant differences between the treatments in the number of branches. The humic acid @ 4 ml l⁻¹ (H_3) gave the maximum branches (7.58) compared with control. The interaction between both factors significantly affected on branches number produced per sapling. The C_3H_3 gave the highest average branch sapling⁻¹ (9.58) compared with (C_1H_1).

Stem diameter (mm): There were significant differences in the stem diameter average recorded in the different types of the cuttings. The highest diameter was shown in the basal cuttings (C_3) which was (12.82 mm), and significantly different from the other cuttings types, while the terminal cuttings (C_1) gave the lowest diameter (10.76 mm) (Table 2). The effect of humic acid was clear when added, so the treatment (H_3) gave the highest average (13.00 mm) which significantly differed with the non addition treatment (H_1) which recoded the lowest average (9.88 mm). Furthermore, the interaction between both studied factors, showed the significant effect, especially for the treatment (C_3H_2) which

Table 1. Some chemical and physical properties of media soil

Mg mmol L ⁻¹	Ca mmol L ⁻¹	K Av. g kg ⁻¹	P Av. g kg ⁻¹	N Av. g kg ⁻¹	O.M g kg ⁻¹	EC ds m ⁻¹	pH
37	43	21.7	17.4	2.56	2.3	1.8	7.8
Texture	Sand g kg ⁻¹	Silt g kg ⁻¹	Clay g kg ⁻¹	HCO ₃ mmol L ⁻¹	SO ₄ mmol L ⁻¹	CO ₃ mmol L ⁻¹	Na mmol L ⁻¹
Clay loam	422.4	210	367.6	4	9.6	Nil	13.1

Table 2. Effect of cutting type and humic acid on vegetative growth traits of fig Waziri cv. saplings

Cutting type	Humic (ml l ⁻¹)	Saplings height (cm)	Number of branches (branches sapling ⁻¹)	Stem diameter (mm)	Sapling leaves area (cm ² sapling)	Dry weight of vegetative system (g)
Terminal cuttings	0	35.79	5.63	9.41	145.5	15.18
	2	46.31	7.51	9.87	148.8	17.44
	4	50.53	6.00	13.00	179.7	21.52
Middle cuttings	0	46.94	6.97	10.09	158.2	16.93
	2	43.75	6.30	11.48	180.8	18.15
	4	56.44	7.16	12.65	175.1	21.62
Basal cuttings	0	41.87	6.14	10.16	183.3	16.70
	2	54.42	7.34	14.94	184.4	20.91
	4	53.30	9.58	13.35	173.9	20.77
LSD (0.05)		7.55	1.74	2.30	17.69	N.S

gave the highest average (14.94 mm) compared with non-humic addition treatment for terminal cuttings (C_1H_1) in which the stem diameter reached the lowest average 9.41 mm (Table 2).

Sapling leaves area (cm^2): The leaf area had significantly differed among the various stem cuttings. The basal cuttings (C_3) gave the highest average (180.5 cm^2) which is not significantly different from the middle cuttings (C_2) but differed from the terminal cuttings (C_1) which registered the lowest leaves area (158.0 cm^2). Additionally, the addition of humic acid resulted in significant differences in this trait, and the treatment (H_3) gave the highest leaves area (176.2 cm^2) which is significantly differed from the non-application treatment (H_1) which recorded the lowest average (162.3 cm^2). The interaction between the two factors showed significant differences among the treatments (Table 2). The treatment (C_3H_2) gave the highest leaves area (184.4 cm^2), while the non-addition of humic acid to the terminal cuttings (C_1H_1) gave the lowest average (145.5 cm^2). The humic acid participates supporting nutritional minerals for chlorophyll biosynthesis and improving roots ability to absorb that minerals (Arancon et al 2006), especially nitrogen, phosphorus and potassium which have important role in chlorophyll synthesis by participating in the composition of proteins and amino-acids and activating enzymes that functionally building them. It's clear that the basal cuttings and humic acid application have a positive role in leaves area increase for saplings.

Dry weight of vegetative system (g): The using of different types of cuttings didn't affect significantly in the dry weight of the vegetative system. While, the humic acid addition significantly affected in this trait, and the treatment (H_3) gave the highest dry weight (21.30 g) compared with the lowest value (16.26 g) recorded at the non-addition treatment (H_1). On the other hand, the interaction between the two studied factors significantly affected this trait (Table 2). the highest mean (21.62 g) recorded by the treatment (C_2H_3) treatment compared with the lowest dry weight of the vegetative system (15.18 g) shown in the treatment (C_1H_1).

Roots length (cm): Root length has significantly differed among cutting types used in the experiment, where the basal and middle cuttings (C_2 and C_3) gave the highest average (41.61 and 40.73cm) with an increase percentage of (17.94 and 15.45%) respectively, from the longest roots in the terminal cuttings (C_1) which recorded the lowest average (35.28 cm). The superiority of basal cuttings on terminal ones in root growth traits could be attributed to be contained more quantity stored nutrients such as carbohydrate, which play the important role in rooting process, especially at the early stages of root formation in the cuttings (Karakurt et al 2009).

Palanisamy and Kumar (1997) also obtained that there's a positive relationship between taken cuttings' diameter and number and length of roots formed, which could be explained for the thicker cuttings have more quantity of carbohydrate that is necessary to form roots (Hartmann et al 2002).The addition of humic acid to saplings also caused significant differences in their root length. The treatment (H_3) gave the highest average (42.14 cm) with an increase percentage of 18.14% compared on the non-humic acid addition (H_1) which gave the lowest average (35.67cm). Furthermore, the both studied factors interaction significantly affected in this trait (Table 3). The treatment (C_2H_3) treatment gave the highest average (45.41 cm), while the non-addition of humic acid treatment to the terminal cuttings (C_1H_1) recorded the lowest average (31.54 cm).

Dry weight of the root system (g): Dry weight of the root system has been significantly affected by the cuttings type used in this experiment, where the middle cuttings (C_2) gave the highest average (8.35 g) compared with the basal and terminal cuttings (C_1 and C_3) which gave the averages (7.88 and 7.61g), respectively. The superiority of basal cuttings on the dry weight of root growth could be attributed to be contained more quantity stored nutrients such as carbohydrate, which play an important role in rooting process, especially at the early stages of root formation in the cuttings (Karakurt et al 2009). The increase in rooting supporting materials such as auxins and other rooting factors in the basal cuttings could be a reason to be led this result (Owais 2010, Rana and Sood 2012). Additionally, the humic acid addition significantly affected the same trait, where the treatment (H_3) gave the highest dry weight of the root system (8.43 g), followed by (H_2), which gave 8.07 g, while the lowest value (7.33 g) recorded in the non-humic acid addition treatment (H_0). The interaction between both studied factors

Table 3. Effect of cutting type and humic acid interaction on root growth traits of figs saplings cv. Waziri

Cutting type	Humic ($ml L^{-1}$)	Roots length (cm)	Dry weight of root system (g)
Terminal cuttings	0	31.54	6.86
	2	35.76	7.01
	4	38.53	8.94
Middle cuttings	0	33.18	7.73
	2	43.58	9.11
	4	45.41	8.22
Basal cuttings	0	42.30	7.40
	2	40.06	8.10
	4	42.47	8.13
LSD (0.05)		5.08	0.98

significantly affected this trait, especially, the treatment (C₂H₂), which gave the highest average (9.11 g). Whereas, the lowest dry weight of the root system appeared in the non-humic with addition for terminal cuttings (C₁H₁), which gave 6.86 g (Table 3). The humic acid has a role in increasing the cation exchange capacity, reducing soil pH and make macro and micro nutrients more available, as well as building a highly efficient root system to absorb nutrients, which positively reflected on plant growth (Eissa et al 2007). Also humic acid participates supporting nutritional minerals for chlorophyll biosynthesis and improving roots ability to absorb those minerals (Arancon et al 2006).

CONCLUSION

Accordingly, the results obtained from this study, it's possible to conclude that the basal cuttings resulted, a positive effect on the studied vegetative and root traits compared on the middle and terminal cuttings. Furthermore, the application of humic acid at (4 ml l⁻¹) concentration improved vegetative and root growth of fig saplings under this study. So, we recommend using basal cutting in fig propagation, as well as, adding humic acid to improve vegetative and root growth traits of fig saplings.

REFERENCES

- Arancon NQ, Clive AE, Stephen L and Robert B 2006. Effects of humic acids from vermin composts on plant growth. *European Journal of Soil and Biology* **42**: 65-69.
- Atee AS and AL-Sahaf FH 2007. Potato production by organic farming; 1-role of organic fertilizer on soil physical properties and microorganism number. *The Iraqi Journal Agricultural Sciences* **38**(4): 36-51.
- Awad EM 2002. Effect of compost and some biofertilizers on growth, yield and quality of potato crop (*Solanum tuberosum* L.). *Journal Agriculture Science* **27**: 5525-5537.
- Bhattacharya S and Nanda KK 1978. Stimulatory effect of purine and pyrimidine bases and their role in the mediation of auxin action through the regulation of carbohydrate metabolism during adventitious root formation in hypocotyl cutting of *Phaseolus mungo*. *Zeitschrift für Pflanzenphysiologie* **88**: 283-293.
- Central System for Statistics and Information Technology (CSSIT) 2018. *Report of Trees Production of Summer Fruits*. Directorate of Agricultural Statics. Ministry of Planning and Development Cooperation. Republic of Iraq.
- Eissa FM, Fathi M and EL-Shall SA 2007. Response of peach and apricot seedling to humic acid treatment under salinity condition. *Journal of Agriculture Science* **32**(5): 3605-3620.
- Eman AA, Abd El-Monerm M, Saleh S and Mostafa EAM 2008. Minimizing the quantity of mineral nitrogen fertilizers on grapevine by using humic acid, organic and biofertilizers. *Res. Journal of Agriculture and Biological Science* **4**(1): 46-50.
- Ferrara G and Brunetti G 2010. Effects of the times of application of a soil humic acid on berry quality of table grape (*Vitis vinifera* L.) cv. Italia. *Spanish Journal Agriculture Research* **8**(3): 817-822.
- Hamooh BT 2004. Cuttings types and IBA concentrations in relation to rooting of stem hardwood cuttings of fig tree (*Ficus carica* L.). *Annals Agriculture Science* **49**(2): 661-669.
- Harrison RD 2005. Figs the university of tropical rain forests. *Bio Science* **55**: 1053-1064.
- Hartmann HT, Kester DE Davies FT and Geneve RL 2002. *Plant Propagation 'Principles and Practices'*. 7th Edition. Prentice Hall. New Jersey.
- Herre EA, Jan KC, Machado CA 2008. Evolutionary ecology of fig leaves extra; Recent pro grass and out staneling Puzles. *Ann Per Ecol Evol Syst* **37**: 438-456.
- Karakurt H, Aslantas R, Ozkan G and Guleryuz M 2009. Effect of indole 3- butyric acid (IBA) plant growth promoting rhizobacteria (PGPR) and carbohydrates on rooting of hardwood cutting of MM106 Apple rootstock. *African Journal of Agriculture Research* **4**(2): 60-64.
- Katkat AV, Gelick H, Turan MA and Asik BB 2009. Effect of soil and foliar application of humic substances on dry weight and mineral nutrients of wheat under calcareous soil conditions. *Australian Journal of Basic and Applied Sciences* **3**(2): 1266-1273.
- Khapare LS, Dahale MH and Bhusari RB 2012. Effect of plant growth regulators on rooting in cuttings of fig (*Ficus carica* L.) cv. Dinker. *Asian science. Hind Institute of Science and Technology India* **7**(1): 25-27.
- Mars M, Chatti KO, Salhi-Hanna A, Trifi M and Marrachi M 2008. Fig cultivation and genetic resources in Tunisia. *An Overview Actuator* **798**: 27-32.
- Taj Al-Deen MM and Al-Barakat HNK 2016. Effect of biofertilizer and humic, fulvic application on availability of some elements in soil. *Al-Muthana Journal of Agriculture Sciences* **4**(2): 1-13.
- Owais SJ 2010. Rooting response of five pomegranate varieties to indole butyric acid concentration and cuttings age. *Pakistan Journal of Biological Science* **13**(2): 51-58.
- Palanisamy K and Kumar P 1997. Effect of position, size of cuttings and environmental factors on adventitious in neem (*Azadirachta indica* A. Juss). *Forest Ecology and Management* **98**: 277-288.
- Pettit RE 2003. Emeritus Associate Professor Texas A & M university, Organic Matter, Humus, Humates Humic Acid, Fulvic Acid and Humin: Their Importance in Soil Fertility and Plant Health. *CTI Research*, **1**, pp.1-17.
- Rana RS and Sood KK 2012. Effect of cutting diameter and hormonal application on the propagation of *Ficus roxburghii* wall through branch cuttings. *Annals of Forest Research* **55**(1): 1-16.
- Reddy KV, Pulla C and Goud PV 2008. Effect of auxins on the rooting of fig (*Ficus carica* L.) hardwood and semi hardwood cuttings. *Indian Journal Agriculture Research* **42**(1): 75-78.
- Ronsted NG, Weiblen D, Cook J Salamin MN Machade CA and Savolainen V 2005. 60 Million years of codivergence in the fig-wasp tymbiosis. *Proceedings of the Royal Society of London* **272**: 2593-2599.
- Schnitzer M, Khan SU 1971. *Humic Substances in the Environment*. Dekker. New York.
- Walaga C 2005. *Organic agriculture in Kenya and Uganda*. www.anancy.net.