

## EFFECT OF ADDITION OF DIFFERENT LEVELS OF GINGER (*ZINGIBER OFFICIALE*) AND VITAMIN E TO THE RATIONS OF AWASSI EWES ON SOME BLOOD TRAITS

Audai Sabah Asker<sup>1\*</sup>, Hebat-Alla A. A. Alhamdani<sup>2</sup> and Adel A. Alhamdani<sup>1</sup>

<sup>1</sup>College of Agriculture, University of Anbar, Anbar, Iraq.

<sup>2</sup>College of Education for Women, University of Anbar, Anbar, Iraq.

\*e-mail: audai.sabah@uoanbar.edu.iq

(Received 23 September 2020, Revised 12 December 2020, Accepted 11 January 2021)

**ABSTRACT :** The study was undertaken to compare the effect of addition of different levels of ginger and vitamin E to the ration cellular and biochemical blood characteristics in Awassi ewes. The experiment was conducted on Awassi ewes, aged between 3-4 years with a mean weight  $37\pm 5$  kg, presented in the private farm in Ramadi, Anbar. The animals were divided into five groups (5 ewes for each group). The first group ( $T_1$ ) not treated serve as a control, second ( $T_2$ ), the third group ( $T_3$ ) and the fourth group ( $T_4$ ) were treated with 10, 15 and 20 gm from ginger three times weekly for each ewe. While, the fifth group ( $T_5$ ) treated with 500mg vitamin E three times weekly for each ewe, blood samples were collected at day 15, 30, 45, 60 of the experiment. The results showed that there was a significant increase in pcv %, Hb %, the numbers of RBCs, WBC, lymphocytes and platelets in  $T_4$  during all the periods. While, the biochemical characteristics of serum showed a high significant value of total protein and globulins for  $T_4$  and  $T_5$  in the most of the periods. Fourth treatment showed significant decrease on the values of glucose, cholesterol and urea at all periods except the value of cholesterol at the first period. The result showed there was no significant difference in the concentration of albumin between different treatment at all periods. It was concluded that the ginger and vitamin E have beneficial effect on blood traits in the ewes.

**Key words :** Ginger, vitamin E, Awassi ewes, blood traits.

**How to cite :** Audai Sabah Asker, Hebat-Alla A. A. Alhamdani and Adel A. Alhamdani (2021) Effect of addition of different levels of ginger (*Zingiber officiale*) and vitamin E to the rations of Awassi ewes on some blood traits. *Biochem. Cell. Arch.* **21**, 727-732. DocID: https://connectjournals.com/03896.2021.21.727

### INTRODUCTION

The most important things for animal physiologist was to keep the animal Hemostasis, which means that all body organs showed normal function that leads to more productive performance. As a results of unsuitable environmental and managemental conditions that the animal suffering from it which may leads to deviation in animal body homeostasis, causing a disturbance in productive performance. In order to correct this deviation, it showed improve a good environmental and managemental conditions. One of these improvement by a researchers was to use herbal additives to diet which considered as antioxidant and antimicrobial or it may have a role to provide some of nutritive elements or it may play a role similar to hormones and enzymes (Lemma and Egza, 2019). The ginger (*Zingiber officinale*) regarded as one of herbal plants have an active effect on biological activity on animal body (Shahrajabian *et al*,

2019). Ginger contained a natural substances accepted by the body without side effects, it improves the digestive enzyme activity such as; Amylase, Lipase and trypsin (srinivagen, 2003; Rahman *et al*, 2016). It is also reducing of intestinal wall irritation, removal of gases and stimulate digestion in cattle (Pakashi, 2003). Ginger acts to increase of bile secretion which is important for fat digestion (Mabey, 1988). On the other hands, ginger reduce the observation of cholesterol and decrease its concentration in the blood serum. Various ginger extracts tend to decrease oxidation of low lipoprotein (LDLP). The ginger contains zingerone, gingerol, and gengeriorol which are a compound having similar activity of vitamin E to prevents oxidation and improved blood characteristics (Mao *et al*, 2019; Anwar *et al*, 2020). This study was aimed to show the effect of different levels of ginger powder and vitamin E on blood characteristics of Awassi ewes.

## MATERIALS AND METHODS

The study was conducted to compare the effect of addition of different levels of ginger (*Zingiber officinal*) with the vitamin E to the ration on cellular and biochemical blood traits in Awassi ewes. The study was carried out on 25 Awassi ewes, aged between 3-4 years, with a mean weight of  $37 \pm 5$  kg, presented in the private farm in Ramadi-Anbar, during a period of two months. The ewes were feed on hay with straw addlbitum with a mean of concentrate 400 gm daily. The ewes were treated with ant parasite drug for external and internal parasite and vaccinated against Brucellosis. The ewes were divided into five groups (5 ewes for each group). The first group ( $T_1$ ) not treated and as a control one. The second ( $T_2$ ), the third group ( $T_3$ ) and the fourth group ( $T_4$ ) were treated with 10, 15, 20 gm from ginger three times weekly for each ewe, respectively, while the fifth group ( $T_5$ ) treated with 500mg from vit. E three times weekly for each ewe. Blood samples were collected via jugular vein at day 15, 30, 45 and 60 of the experiment to measure the following parameters: complete blood picture; measurement of red blood cells count according to lewis and daue (1974), white blood cells (WBC) according to Joha and Lewis (1984), packed cell volume (PCV) according to Lewis and Daue (1974), Hemoglobin (Hb%), blood platelets according to Schalm *et al* (1975) and the lymphocytic cells according to Seiverd (1973). Parameters of some blood biochemistry: The measurement of cholesterol according to Tietz (1995) and glucose according to Kopper (1975), measurement of total protein and albumin according to the analytical Kits manufactured by biolab Co. France and according to the guide of the kit. Globulins were measured according to the method of Freeman and Wotton (1974), which described by Bishop *et al* (2000). The urea was measured according to Freem and Wotton (1974). Statistical analysis was done using one-way analysis and the use of SAS program (SAS, 2002) 9.1. Also multiple range test were applied (Duncan, 1955). 5% level of significant were use between different values ( $P \leq 0.05$ ).

## RESULTS AND DISCUSSION

The results showed in Table 1 that at the first period (day 15). There was a significant increase ( $P \leq 0.05$ ) in all treated groups as a compared with the control one in the cellular traits of the blood (PVC, HB, RBC, WBC, lymphocytes and blood platelets) numbers, Treatment number four ( $T_4$ ) showed the highest significant values ( $P \leq 0.05$ ) for these traits. The concentration of serum total protein and globulin showed significant increase ( $P \leq 0.05$ ) in all treated groups as a compared with Control one, but  $T_4$  and  $T_5$  showed a high significant value

**Table 1** : Effect of ginger in blood tests of ewes at 15 days of experiment.

Tests	Treatments					P-value
	$T_1$	$T_2$	$T_3$	$T_4$	$T_5$	
PCV%	$22.4 \pm 0.278^d$	$24.3 \pm 0.205c$	$27.8 \pm 0.379b$	$31.3 \pm 0.299a$	$30.6 \pm 0.405a$	0.0001
Hb%	$7.48 \pm 0.146c$	$8.45 \pm 0.101b$	$8.34 \pm 0.088b$	$10.8 \pm 0.107a$	$10.5 \pm 0.132a$	0.0001
RBC (million/ml <sup>3</sup> )	$6.36 \pm 0.260d$	$8.18 \pm 0.093c$	$8.96 \pm 0.187bc$	$10.73 \pm 0.480a$	$9.84 \pm 0.554ab$	0.0001
WBC (thousand/ml <sup>3</sup> )	$13.31 \pm 0.681d$	$20.68 \pm 0.582c$	$26.44 \pm 0.766b$	$31.22 \pm 0.720a$	$27.17 \pm 1.06b$	0.0001
Lymphocyte (thousand/ml <sup>3</sup> )	$5.98 \pm 0.214e$	$20.11 \pm 0.897d$	$26.01 \pm 0.536c$	$31.05 \pm 0.393a$	$27.72 \pm 0.461b$	0.0001
Platelets (thousand/ml <sup>3</sup> )	$434 \pm 12.0c$	$504 \pm 5.03b$	$515 \pm 2.23b$	$551 \pm 4.00a$	$549 \pm 4.07a$	0.0001
Cholesterol (mg/dl)	$17.1 \pm 0.366$	$17.1 \pm 0.335$	$16.8 \pm 0.216$	$16.4 \pm 0.481$	$17.4 \pm 0.453$	N.S.**
Glucose (mg/dl)	$76.1 \pm 0.628a$	$67.9 \pm 0.476b$	$64.9 \pm 1.13c$	$54.4 \pm 1.42d$	$68.9 \pm 0.855b$	0.0001
Urea (mg/dl)	$67.4 \pm 1.73a$	$66.1 \pm 1.15a$	$53.6 \pm 1.22b$	$51.0 \pm 0.359b$	$67.8 \pm 1.26a$	0.0001
Protein (gm/dl)	$4.99 \pm 0.071c$	$5.34 \pm 0.040b$	$5.46 \pm 0.074b$	$6.05 \pm 0.096a$	$6.09 \pm 0.096a$	0.0001
Albumin (gm/dl)	$2.94 \pm 0.099$	$2.92 \pm 0.090$	$2.97 \pm 0.105$	$2.86 \pm 0.121$	$3.00 \pm 0.047$	N.S.
Globulin (gm/dl)	$2.05 \pm 0.133c$	$2.42 \pm 0.093b$	$2.48 \pm 0.121b$	$2.99 \pm 0.085a$	$3.09 \pm 0.140a$	0.0001

\* Means  $\pm$  Standard Error. \*\* N.S.: Non-Significant. a, b, c: means in the same Rows with different superscripts differ significantly at probability value ( $P \leq 0.05$ ).

**Table 2 :** Effect of ginger in blood tests of ewes at 30 days of experiment.

Tests	Treatments					P-value
	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>	T <sub>5</sub>	
PCV%	22.9 ± 0.132 <sup>d</sup>	24.8 ± 0.272 <sup>c</sup>	28.2 ± 0.467 <sup>b</sup>	32.1 ± 0.271 <sup>a</sup>	31.4 ± 0.202 <sup>a</sup>	0.0001
Hb%	7.90 ± 0.171 <sup>c</sup>	8.19 ± 0.084 <sup>bc</sup>	8.45 ± 0.075 <sup>b</sup>	10.50 ± 0.144 <sup>a</sup>	10.38 ± 0.111 <sup>a</sup>	0.0001
RBC (million/ml <sup>3</sup> )	6.29 ± 0.145 <sup>c</sup>	8.50 ± 0.217 <sup>b</sup>	8.50 ± 0.269 <sup>b</sup>	11.17 ± 0.348 <sup>a</sup>	10.65 ± 0.404 <sup>a</sup>	0.0001
WBC (thousand/ml <sup>3</sup> )	14.29 ± 0.936 <sup>d</sup>	22.58 ± 0.472 <sup>c</sup>	26.19 ± 0.551 <sup>b</sup>	31.36 ± 0.365 <sup>a</sup>	25.92 ± 0.521 <sup>b</sup>	0.0001
Lymphocyte (thousand/ml <sup>3</sup> )	5.84 ± 0.129 <sup>d</sup>	20.57 ± 0.696 <sup>c</sup>	26.24 ± 0.599 <sup>b</sup>	30.95 ± 0.592 <sup>a</sup>	27.71 ± 0.629 <sup>b</sup>	0.0001
Platelets (thousand/ml <sup>3</sup> )	456 ± 6.78 <sup>c</sup>	510 ± 5.49 <sup>b</sup>	515 ± 1.80 <sup>b</sup>	549 ± 5.09 <sup>a</sup>	552 ± 3.72 <sup>a</sup>	0.0001
Cholesterol (mg/dl)	17.0 ± 0.342 <sup>a</sup>	14.4 ± 0.314 <sup>b</sup>	14.4 ± 0.633 <sup>b</sup>	12.3 ± 0.240 <sup>c</sup>	15.0 ± 0.597 <sup>b</sup>	0.0001
Glucose (mg/dl)	77.3 ± 1.09 <sup>a</sup>	66.9 ± 0.538 <sup>b</sup>	66.6 ± 0.580 <sup>b</sup>	53.4 ± 0.398 <sup>c</sup>	68.0 ± 0.819 <sup>b</sup>	0.0001
Urea (mg/dl)	64.8 ± 1.09 <sup>a</sup>	58.6 ± 0.894 <sup>b</sup>	51.0 ± 0.775 <sup>c</sup>	50.8 ± 0.259 <sup>c</sup>	67.0 ± 0.544 <sup>a</sup>	0.0001
Protein (gm/dl)	5.03 ± 0.053 <sup>c</sup>	5.48 ± 0.058 <sup>b</sup>	5.48 ± 0.086 <sup>b</sup>	6.12 ± 0.123 <sup>a</sup>	6.09 ± 0.086 <sup>a</sup>	0.0001
Albumin (gm/dl)	3.09 ± 0.126	2.99 ± 0.152	2.84 ± 0.205	2.97 ± 0.153	2.91 ± 0.082	N.S.**
Globulin (gm/dl)	1.93 ± 0.159 <sup>c</sup>	2.48 ± 0.163 <sup>b</sup>	2.63 ± 0.157 <sup>b</sup>	3.07 ± 0.071 <sup>a</sup>	3.16 ± 0.136 <sup>a</sup>	0.0001

\* Means ± Standard Error. \*\* N.S.: Non-Significant. a, b, c: means in the same Rows with different superscripts differ significantly at probability value (P≤0.05).

(P≤0.05) as compared with other treatment. There was a significant decrease in the glucose and urea concentration (P≤0.05) in T<sub>3</sub> and T<sub>4</sub> as compared with other treatment. T<sub>4</sub> showed the lowest significant (P≤0.05) in the concentration of glucose. There was no significant difference between treated group in the concentration of cholesterol and albumin.

The results also showed at the second period (day 30) (Table 2), that there was a significant increase (P≤0.05) for the T<sub>4</sub> and T<sub>5</sub> as compared other with other treated group in the percentage of PCV, Hb, RBC, blood platelets and the concentrations of the total protein and globulin. T<sub>4</sub> showed significant increases (P≤0.05) in WBC and lymphocytic count while T<sub>4</sub> showed a significant decrease (P≤0.05) in the conc. Of cholesterol and glucose compared with other treated groups. The T<sub>3</sub> and T<sub>4</sub> showed significant decrease (P≤0.05) in the conc. Of urea as compared with other treatments, while there were no differences in the conc. of albumin in all treatments.

The results of the third period (Day 45) (Table 3) showed a significant increase (P≤0.05) in T<sub>4</sub> and T<sub>5</sub> as compared with other treatments in the percentage of HB, RBC, Platelets no. and total protein. The result also showed a significant increase (P≤0.05) in T<sub>4</sub> in PVC as compared with others, while the conc. of globulin showed an increase in the concentration significantly (P≤0.05) in T<sub>5</sub> as compared with T<sub>1</sub>, T<sub>2</sub> and T<sub>3</sub>, T<sub>4</sub> showed an increase in its value as compared with T<sub>1</sub> and T<sub>3</sub>. The conc. of cholesterol and glucose showed significant decrease (P≤0.05) in the T<sub>4</sub> as compared with others. The conc. of albumin showed no difference between the four treated groups.

The results of the fourth period (day 60) showed in Table 4, this period were not differing from that of previous periods it showed no significant increase (P≤0.05) in percentage of PCV, Hb and platelets counts in T<sub>4</sub> and T<sub>5</sub> as compared with other treatments. T<sub>4</sub> showed a significant increase (P≤0.05) in RBC, WBC and lymphocytic cells as compared with others. There was a significant increase (P≤0.05) in total protein at T<sub>4</sub> as compared with the first three treatments. It was also showed T<sub>4</sub> and T<sub>5</sub> a significant increase (P≤0.05) in the conc. of globulin as compared with T<sub>1</sub> and T<sub>2</sub>. While, T<sub>4</sub> showed significant decrease (P≤0.05) in the conc. of cholesterol, glucose and urea as compared with other treatment. The conc. of albumin showed no difference in other periods in all treatment.

The significant difference observed in PCV, Hb, blood platelets and RBC values, which produced from addition

of various levels of ginger and vit. E as compared with the control, might be due to the role of ginger to stimulate apatite and improve digestion of food material consumed, that leads to beneficial effect from food elements especially vitamins and proteins that plays a role in the synthesis of RBCs that effects of another parameters of blood (PVC and Hb %) and the blood platelets (Hung, 1991; Al-Saidia, 2010). Ginger contains Fe (Mabey, 1988), which is essential for the synthesis of RBCs or it might be due to the presence of antioxidant in ginger such as gingerol, shagaols and phenyl kiton and also due to the action of vit. E that protect the body cells from free radicals which include secretion cells that enhance the synthesis of RBCs such as ESF, which secreted from the kidney so increase RBC and Hb when a positive correlation presence between them. The antioxidant that mentioned and vit. E protect blood platelets and increase its synthesis and life. The active substances present in ginger and vit. E play a role in protect the organ produced white blood cells such as thymus gland and bone marrow from the toxic effect of free radicals although it protects WBC and prevents the effects of free radicals and increase its life that leads to increase their number (Hadi, 2009). The improvement of immune response as a result of addition of ginger and vit. E might be due to the content of ginger of multi active substances that act synergistically to improve blood cells characteristics. The significant improvement in biochemical blood traits understudy except in conc. of albumin might be due to addition of ginger and vit. E to protect body cells such as liver cells and Beta-cells of pancreas from the side effect or advirse effect of free radicals produced from biological activity of the body and the active substances present in ginger tend to reduce lipid levels (Al-Janabi, 2004; Sabeg, 2010) or it might be due to content of ginger of an active substances such gingerol that prevents inflammation in the body and protect Beta-cells of pancreas that secret insulin leads to decrease in glucose conc. In the blood which stimulates cells receptors to facilitate the passage of glucose into the body cells to produce energy essential for maintenance of biological activity in the body (Sekua *et al*, 2004; Ashokkumar *et al*, 2020). The decrease in cholesterol level might be due to addition of ginger that have a biological active substance inhibits the yeast (HMG COA red), which limits biological maturation of cholesterol (Sabour *et al*, 2012; Noaman, 2018). The significant increase in protein conc. might be due to the role vit. E and ginger on body cells to protect epithelial line of the digestive system that increase absorpction of amino acids, which given with the ration and provide a good condition for microbial digestion of food substance consumed that

**Table 3 :** Effect of ginger in blood tests of ewes at 45 days of experiment

Tests	Treatments					P-value
	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>	T <sub>5</sub>	
PCV%	22.1 ± 0.562 <sup>e</sup>	24.8 ± 0.275 <sup>d</sup>	28.6 ± 0.244 <sup>c</sup>	31.6 ± 0.317 <sup>a</sup>	30.3 ± 0.500 <sup>b</sup>	0.0001
Hb%	7.57 ± 0.165 <sup>c</sup>	8.52 ± 0.066 <sup>b</sup>	8.77 ± 0.040 <sup>b</sup>	10.60 ± 0.151 <sup>a</sup>	10.72 ± 0.131 <sup>a</sup>	0.0001
RBC (million/ml <sup>3</sup> )	6.06 ± 0.161 <sup>c</sup>	8.54 ± 0.204 <sup>b</sup>	8.84 ± 0.062 <sup>b</sup>	10.92 ± 0.451 <sup>a</sup>	10.41 ± 0.585 <sup>a</sup>	0.0001
WBC (thousand/ml <sup>3</sup> )	15.68 ± 0.413 <sup>c</sup>	23.71 ± 0.936 <sup>b</sup>	26.16 ± 1.16 <sup>b</sup>	31.11 ± 0.439 <sup>a</sup>	25.29 ± 0.899 <sup>b</sup>	0.0001
Lymphocyte (thousand/ml <sup>3</sup> )	6.31 ± 0.262 <sup>d</sup>	20.98 ± 0.645 <sup>c</sup>	26.69 ± 0.727 <sup>b</sup>	30.99 ± 0.401 <sup>a</sup>	26.62 ± 0.756 <sup>b</sup>	0.0001
Platelets (thousand/ml <sup>3</sup> )	452 ± 5.46 <sup>c</sup>	511 ± 2.22 <sup>b</sup>	516 ± 4.30 <sup>b</sup>	555 ± 4.47 <sup>a</sup>	554 ± 6.96 <sup>a</sup>	0.0001
Cholesterol (mg/dl)	17.5 ± 0.453 <sup>a</sup>	14.3 ± 0.522 <sup>b</sup>	13.9 ± 0.509 <sup>b</sup>	12.0 ± 0.292 <sup>c</sup>	15.2 ± 0.413 <sup>b</sup>	0.0001
Glucose (mg/dl)	79.0 ± 0.962 <sup>a</sup>	64.7 ± 1.03 <sup>b</sup>	66.0 ± 0.986 <sup>b</sup>	53.0 ± 0.630 <sup>c</sup>	66.0 ± 0.493 <sup>b</sup>	0.0001
Urea (mg/dl)	66.5 ± 1.15 <sup>a</sup>	54.6 ± 0.900 <sup>b</sup>	51.5 ± 1.45 <sup>bc</sup>	48.2 ± 2.38 <sup>c</sup>	65.7 ± 1.15 <sup>a</sup>	0.0001
Protein (gm/dl)	4.99 ± 0.055 <sup>c</sup>	5.56 ± 0.143 <sup>b</sup>	5.52 ± 0.106 <sup>b</sup>	6.13 ± 0.136 <sup>a</sup>	6.19 ± 0.085 <sup>a</sup>	0.0001
Albumin (gm/dl)	2.99 ± 0.096	2.97 ± 0.109	2.98 ± 0.125	3.15 ± 0.074	2.86 ± 0.121	N.S.**
Globulin (gm/dl)	2.00 ± 0.120 <sup>d</sup>	2.59 ± 0.181 <sup>bc</sup>	2.54 ± 0.101 <sup>c</sup>	2.98 ± 0.152 <sup>ab</sup>	3.33 ± 0.131 <sup>a</sup>	0.0001

\* Means ± Standard Error. \*\* N.S.: Non-Significant. a, b, c: means in the same Rows with different superscripts differ significantly at probability value (P≤0.05).

Table 4 : Effect of ginger in blood tests of ewes at 60 days of experiment.

Tests	Treatments					P-value
	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>	T <sub>5</sub>	
PCV%	22.3 ± 0.204 <sup>d</sup>	24.7 ± 0.664 <sup>c</sup>	28.7 ± 0.366 <sup>b</sup>	31.8 ± 0.321 <sup>a</sup>	31.4 ± 0.192 <sup>a</sup>	0.0001
Hb%	7.40 ± 0.201 <sup>c</sup>	8.42 ± 0.260 <sup>b</sup>	8.90 ± 0.070 <sup>b</sup>	10.92 ± 0.267 <sup>a</sup>	10.68 ± 0.182 <sup>a</sup>	0.0001
RBC (million/ml <sup>3</sup> )	6.47 ± 0.158 <sup>d</sup>	9.00 ± 0.100 <sup>c</sup>	9.09 ± 0.152 <sup>c</sup>	11.72 ± 0.406 <sup>a</sup>	9.92 ± 0.370 <sup>b</sup>	0.0001
WBC (thousand/ml <sup>3</sup> )	14.33 ± 1.06 <sup>d</sup>	24.19 ± 0.535 <sup>c</sup>	26.51 ± 0.714 <sup>b</sup>	31.27 ± 0.350 <sup>a</sup>	26.75 ± 0.798 <sup>b</sup>	0.0001
Lymphocyte (thousand/ml <sup>3</sup> )	6.31 ± 0.279 <sup>d</sup>	22.55 ± 0.279 <sup>c</sup>	26.76 ± 0.207 <sup>b</sup>	30.69 ± 0.244 <sup>a</sup>	26.18 ± 0.657 <sup>b</sup>	0.0001
Platelets (thousand/ml <sup>3</sup> )	440 ± 7.07 <sup>c</sup>	511 ± 3.67 <sup>b</sup>	518 ± 1.22 <sup>b</sup>	560 ± 6.69 <sup>a</sup>	555 ± 10.0 <sup>a</sup>	0.0001
Cholesterol (mg/dl)	18.1 ± 0.602 <sup>a</sup>	13.4 ± 0.548 <sup>b</sup>	14.1 ± 0.597 <sup>b</sup>	11.5 ± 0.318 <sup>c</sup>	13.8 ± 0.539 <sup>b</sup>	0.0001
Glucose (mg/dl)	76.7 ± 0.810 <sup>a</sup>	66.0 ± 0.578 <sup>b</sup>	66.9 ± 0.679 <sup>b</sup>	53.6 ± 0.700 <sup>c</sup>	66.5 ± 1.13 <sup>b</sup>	0.0001
Urea (mg/dl)	66.5 ± 1.13 <sup>a</sup>	53.5 ± 0.774 <sup>b</sup>	50.7 ± 1.03 <sup>b</sup>	41.8 ± 0.902 <sup>c</sup>	66.1 ± 1.27 <sup>a</sup>	0.0001
Protein (gm/dl)	4.88 ± 0.125 <sup>d</sup>	5.48 ± 0.086 <sup>c</sup>	5.64 ± 0.146 <sup>bc</sup>	6.08 ± 0.093 <sup>a</sup>	5.93 ± 0.145 <sup>ab</sup>	0.0001
Albumin (gm/dl)	3.24 ± 0.118	3.22 ± 0.073	3.02 ± 0.060	2.97 ± 0.051	3.04 ± 0.061	N.S.**
Globulin (gm/dl)	1.63 ± 0.213 <sup>c</sup>	2.26 ± 0.132 <sup>b</sup>	2.62 ± 0.201 <sup>ab</sup>	3.11 ± 0.081 <sup>a</sup>	2.88 ± 0.163 <sup>a</sup>	0.0001

\* Means ± Standard Error. \*\* N.S.: Non-Significant. a, b, c: means in the same Rows with different superscripts differ significantly at probability value (P≤0.05).

leads to positive balance of nitrogen to the ewes (Grzanna *et al*, 2005). The significant improvement in globulin level as a result of addition of different levels of ginger and vit. E might be due to the role of active substances in ginger and the role of vit. E to improve immune response which explained by increase in the numbers of WBC resulted from additives that remove the free radicals and protect the main immune organs (Kinalski *et al*, 2000). While no presence of significant difference in conc. of albumin might be due to good management of ewes, with no infections with diseases especially liver disease. The protection of the liver from disease plays an important role in keeping the conc. of albumin constant. The liver considered the main organs synthesized albumin. The positive significant difference in urea conc. In blood serum might be due to addition on ginger that play role in protein synthesis and protect the kidneys for their functions. It was concluded from this study that addition of ginger and vit. E have apposite effects of blood traits in ewes.

## REFERENCES

- Aljanabi O S I (2007) Possible improvement of reproductive criteria with the use of ginger root after the use of cimetidine in mice male. *Master Thesis*, College of Veterinary Medicine, Baghdad University.
- Al-Saidya A M A (2010) Study of pathological and anti-hyperlipidemic effects of ginger *Zingiber officinalae* in rats exposed to oxidative stress. *Iraqi J. Vet. Sci.* **24**(2), 103-108.
- Anwar S, Almatroudi A, Allemailem K S, Joseph R J, Alikhan A and Rahmani A H (2020) Protective effects of Ginger extract against glycation and oxidative stress-induced health complications: An *In vitro* study. *J. Processes* **8**, 468.
- Ashokkumar K, Murugan M, Dhanya M K, Sathyan T, Raj and Mathews N (2020) Chapter 5. Traditional uses phytochemistry, and pharmacological properties of *Zingiber officinale* essential oil and extracts. In: *Ethnopharmacological Investigation of Indian Spices*. DOI: 10.4018/978-1-7998-2524-1.ch005.
- Bishop M L, Dube-Engelkirk J L and Fody E P (2000) Clinical chemistry principles, correlations, procedures. 4<sup>th</sup> ed. J.B. Lippincott Williams and Wilkins. Philadelphia p (405-416).
- Dacie J V and Lewis S M (1974) *Practical Hematology*. 5<sup>th</sup> ed. The English Language Book Soc., London.
- Grzanna R, Lindmik L and Frondoza G G (2005) Ginger - an herbal medicinal. product. with bread anti - inflammatory actions. *J. Med. Food* **8**(2), 125-132.
- Hadi L E (2009) The Synergistic Effect of *Zingiber officinale* Roots and Vitamin E on some Reproductivity and Physiological Traits in Rations of Kids Iraqi Black Goat. *Master Thesis*, College of Veterinary Medicine, Baghdad University.
- Hung Q, Iwamoto M, Aoki S, TaMata N K and Yamahara J (1991) Anti-5-hydroxytryptamine effect of galano lactone diterpenoid isolated from ginger. *Chem-pharm. Bull.* **39**, 397-399.
- Hymoor S, Alnoory A S and Amreen A (2012) Investigation the Effectiveness of Aqueous Extracts of Ginger *Zingiber officinale* Roscoe on the Levels of Total Cholesterol and Lipoproteins in Rats. *Damascus Univ. J. Health Sci.* **28**(2), 433-443.

- John Sir and Lewis S M (1984) Basic Hematological techniques. Practical hematology, 6<sup>th</sup> ed. 27-45.
- Kinalski M, Sledziewsk A and Telejko B (2000) Lipid peroxidation and scavenging enzyme activity in streptozotocin – induced diabetes. *Acta Diabeto.* **1.37**, 179.
- Kopper AJ (1975) Determination of serum glucose. Analytic chemistry. Kaplan A *et al.* The C.V. Mosby Co. St Louis. Toronto, Princeton. 1316-1324.
- Lemma T S and Egza T F (2019) Extraction and characterization of essential oil from ginger rhizome collected from Arba minch market. *Int. J. Engineering Trends and Technol.* **67**(12), 2231-5381.
- Mabey R A (1988) The new age herbalist Fireside Book, Simon and Schuster INC, New York, P. 128.
- Mao Q Q, Xu X Y, Cao S Y, Gan R Y, Corke H, Beta T and Li H B (2019) Bioactive compounds and Bioactive of Ginger (*Zingiber officiale* Roscoe). *J. Foods* **8**, 185.
- Noaman A I (2018) Effect of two level of ginger (*Zingiber officiale*) powder roots in diet on the productive performance and some blood biochemical parameters of Awassi ewe lambs. *J. Annals Agric. Sci., Moshtohor* **56**(1), 123-126.
- Pakrashi S C and Pakrashi A (2003) *Ginger*. vedamsd, New Delhi.
- Rahman S U, Adhikari A, Ismail M, Shaha M R, Khurram M, Shahid M, Ali F, Haseeb A, Akbar F and Iriti M (2016) Beneficial effects of *Trillium govanianum* Rhizomes in pain and inflammation. *Molecules* **21**, 1095.
- Sabiq K K A (2010) The effect of a ginger-based diet program on the treatment of dyslipidemia. *Master Thesis*, Umm Al Qura University, Mecca.
- Schalm O W, Jain N C and Carro E J (1975) *Vet Hematol.* **3**<sup>rd</sup> Ed. W. Lea and Febiger, Philadelphia, USA.
- Seiverd C E (1973) *Hematology for medical Technologies.* 4<sup>th</sup> ed. Lea and Febiger. Philadelphia. p:117-120.
- Sekiya K, Othani A and Kusano S (2004) Enhancement of insulin sensitivity in adipocytes by ginger. *Bio factors* **22**, 153-156.
- Shahrajabian M H, Sun W and Cheng Q (2019) Clinical aspects and health benefits of ginger (*Zingiber officinale*) in both traditional Chinese medicine and modern industry. *Acta Agriculturae Scandinavica*, section B- *Soil & plant science*, Issn:0906-4710, <https://doi.org/10.1080/09064710.2019.1606930>.
- Srinivasan V, Hamza S, Kyishnamurthy K S and Thankamino C K (2003) Threshold level of soil zinc for optimum production of ginger (*Zingiber officinal* Rose). In: *National seminar on new perspectives in spices medical and aromatic plants*, 69-70.
- Tietz N W (1995) *Clinical guide to laboratory Test.* 3<sup>rd</sup> ed. W.B. Saunders company, Philadelphia, USA.
- Woottone I D P (1974) *Micro analysis in medical biochemistry.* 5<sup>th</sup> ed. Willmer Brothers Limited, Birkenhead, UK.
- Wotton I D P and Freeman H (1974) *Microanalysis in medical biochemistry.* 4<sup>th</sup> ed. Churchill Livingstone, London.