

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/350193729>

# GC–Mass Spectrometry Analysis of Iraqi Moringa Oleifera Seeds Extract

Article in *Indian Journal of Forensic Medicine & Toxicology* · March 2021

CITATIONS

0

READS

83

4 authors, including:



**Abid Thaker**

University of Anbar

22 PUBLICATIONS 72 CITATIONS

[SEE PROFILE](#)



**Asmiet Ramizy**

University of Anbar

84 PUBLICATIONS 521 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Design and Optimization of Nanoparticles Germanium pn junction light-emitting diode - Theoretical study [View project](#)



Nanostructured thin films prepared via pulsed laser ablation in liquid for biological applications [View project](#)

# GC-Mass Spectrometry Analysis of Iraqi *Moringa Oleifera* Seeds Extract

Fiham Jassim AL-Obaidi<sup>1</sup>, Abid A. Thaker<sup>2</sup>, Asmiet Ramizy<sup>3</sup>

<sup>1</sup> MSc. Student, <sup>2</sup> Prof. Department of Biology, college of Science, University of Anbar, <sup>3</sup> Prof., Department of Physics, college of Science, University of Anbar

## Abstract

*Moringa oleifera* is an important traditional medicinal plant used for the treatment of many health problems. It is widely cultivated across the world but it has not well known or well-studied in Iraq. Therefore, the aim of this study is to identify the bioactive compounds present in acetone:methanol (1:1) seeds extract of Iraqi *Moringa oleifera* by using GC-Mass Spectrophotometry.

The results of this analysis revealed 41 compounds which include terpenes ( cis-beta-Ocimene 0.97%, Cedrene 0.20%, Cedrenol 1.40%, Verrucarol 0.29 , Dehydroxy-isocalamendiol 0.09% , beta-Myrcene 0.14% , 3-Thujene 0.06%, Ocimene 0.72% , Sabinene 0.37% Aromadendrene 1.54%) , phenols (0.34%, Eugenol 0.49%, cis-Isoeugenol 1.21%, Methoxyeugenol 0.26%, Ferrocene 0.57%, Phenol, 2-methoxy-4-(1-propenyl)-, (Z) 0.34 , Oxabicyclo[3.3.0]octan-2-one -7-neopentylidene) 0.61% ) , fatty acid methyl esters ( n-Tridecanoic acid 0.15% , Undecanoic acid 0.09% , Pentadecanoic acid 34.43%, 4-Hexadienal 1.52% , Dodecanoic acid 11.02%, cis-Vaccenic acid 21.39%, Hexadecenoic acid 0.15% , Eicosanoic acid 0.03%), acid (myristicin, 7.8%), Phenyl propanoid (Asarone, 4.99%), Antioxidant (-tert-Butyl-4-hydroxyanisole, 1.38%), Drug (Berbine, 2,3,9,10-tetramethoxy, 0.73%), Drug (Thiazolo, 0.63%), aldehyde (Trimethylsilyl vanillin, 0.61%), hormone (Octenoic acid, 0.46%), Amine (ethyl ester, 0.42%), Carboxylic acid (Dimethylandroster-5-en-3-one, 0.28%), Terpenoid (Lavandulol, 0.19%), Heterocyclic compound (Buten-3-one, 1-(2-carboxy-4,4-dimethylcyclobutenyl, 0.18%), Volatile oil (Butanoic acid, 0.16%), Fatty alcohol (1-Heptatriacontanol, 0.12%), Hydroxy carbon-alkene-(Carene, 0.11%).

It can be concluded from this study that the seeds are considered a rich source of bioactive compounds.

**Keywords:** GC-MS analysis, Bioactive compounds, Methanol:Acetone Extract, Iraqi *Moringa oleifera*.

## Introduction

Medical plants are commonly used as a source of medicinal products or against the toxicity of xenobiotics (1,2,3). World Health Organization reports that 4 billion people (80 percent of the world's population) use herbal medicinal goods for most of the primary health care (4).

*Moringa oleifera* is among 13 species of *Moringa* trees that grow in different countries around the world (5).

Seeds of *M. oleifera* tree are useful for medicinal, practical food preparations, water purification, and biodiesel production. (6). *M. oleifera* seeds are a good source of proteins, lipids, vitamins, minerals, micronutrients, and bioactive phytochemicals (7,8,9). The aqueous seeds extract of *M. oleifera* inhibit bacterial and fungal growth (10). The methanol: acetone seed extract of Iraqi *M. oleifera* is found to reduce the toxicity of nanoparticles of lead in the rat (Unpublished results).

This study aims to estimate the compounds found in methanol: acetone seeds extract of Iraqi *M. oleifera* by using GC mass spectroscopy.

---

**Corresponding author:**  
Fiham Jassim AL-Obaidi  
drabidthaker@gmail.com

## Material and Methods

### Plant collection

The ripe dried pods of *M. oleifera* were collected from the trees planted in National Iraqi herbarium (in Baghdad), Iraq, and was identified by a taxonomist in the Desert center, University of Anbar, Iraq.

### Preparation of samples

The seeds were removed from dried pods, dehusked, kept at shade ambient temperature (45°C) to remove any possible moisture. The seeds were later milled to obtain seed powder using a blender and stored it in the refrigerator in airtight containers for analysis.

### Preparation of extract

The dried powdered seeds (300gms) were extracted using Soxhlet and thimble with 1000 ml of solvent (methanol: Acetone 1:1) for 48 hours. The extract was then evaporated with a rotary evaporator. The dried extract was stored at 4°C in a sterile container.

### GC-MS analysis

The chromatograph used was Shimadzu GC-2010 plus-Japan in the Iraqi ministry of sciences and technology; Column: ZB-5MS Capillary Column (30 m x 0.25 mm, I.D. 0.25 µm), Carrier Gas: UHP Helium, Injection Temperature: 280.00°C, Detector Temperature: 280.00 °C, Injection Mode: Split, Flow Control Mode: Pressure, Injector Pressure: 100.0 kPa, Total Flow: 47.3 µl/min, Column Flow: 1.43 µl/min, Linear Velocity: 44.1 cm/min, Injection Volume: 1 µL, Run Time: 35 minutes. A freshly prepared sample (10mg/ml) was used.

## Results and Discussion

GC-MS is an important device used for the analysis of bioactive chemical constituents according to their mass to charge ratio.

Forty one bioactive compounds were identified in seeds extract (methanol:acetone 1:1) of Iraqi *M. oleifera* by GC-MS analysis and detailed in Figure 1., the compounds included terpenes ( cis-beta-Ocimene 0.97%, Cedrene 0.20%, Cedrenol 1.40%, Verrucarol 0.29, Dehydroxy-isocalamendiol 0.09% , beta-Myrcene 0.14% , 3-Thujene 0.06%, Ocimene 0.72% , Sabinene 0.37% Aromadendrene 1.54%) , phenols (0.34%,

Eugenol 0.49%, cis-Isoeugenol 1.21%, Methoxyeugenol 0.26%, Ferrocene 0.57%, Phenol, 2-methoxy-4-(1-propenyl)-, (Z) 0.34 , Oxabicyclo[3.3.0]octan-2-one -7-neopentylidene) 0.61% ) , fatty acid methyl esters ( n-Tridecanoic acid 0.15% , Undecanoic acid 0.09% , Pentadecanoic acid 34.43%, 4-Hexadienal 1.52% , Dodecanoic acid 11.02% , cis-Vaccenic acid 21.39%, Hexadecenoic acid 0.15% , Eicosanoic acid 0.03%), acid (myristicin, 7.8%), Phenyl propanoid (Asarone, 4.99%), Antioxidant (-tert-Butyl-4-hydroxyanisole, 1.38%), Drug (Berbine, 2,3,9,10-tetramethoxy, 0.73%), Drug (Thiazolo, 0.63%), aldehyde (Trimethylsilyl vanillin, 0.61%), hormone (Octenoic acid, 0.46%), Amine (ethyl ester, 0.42%), Carboxylic acid (Dimethylandroster-5-en-3-one, 0.28%), Terpenoid (Lavandulol, 0.19%), Heterocyclic compound (Buten-3-one, 1-(2-carboxy-4,4-dimethylcyclobutenyl, 0.18%), Volatile oil (Butanoic acid, 0.16%), Fatty alcohol (1-Heptatriacontanol, 0.12%), Hydroxy carbon-alkene-(Carene, 0.11%).

The following examples revealed the biological activity of each of the above compounds. Myristicin was found to act as prophylactic effects against colitis in mice, the myristicin treatment before the induction of colitis significantly changed the colonic oxidative stress by elevating the antioxidant enzymatic activities, and reducing the lipid peroxidation<sup>(11)</sup>.


Asarone may be developed to a therapeutic agent to manage cognitive weakening related to conditions such as Alzheimer<sup>(12)</sup> and increase of chemosensitivity by preventing tumor glycolysis in gastric cancer<sup>(13)</sup>.

Methoxyeugenol has anticancer activity and antimicrobial activity<sup>(14)</sup>. Pentadecanoic acid, suppresses the stemness of mcf-7/sc human breast cancer stem-like cells through JAK2/STAT3 signaling<sup>(15)</sup>. Dodecanoic acid, may elicit apoptosis in certain cancer cells<sup>(16)</sup> and exerted antiproliferative activity in diverse types of tumor cells<sup>(17)</sup>. When fed vaccenic acid over 16 weeks, rats exhibited lowered total cholesterol, lowered LDL cholesterol, and lower triglycerides levels<sup>(18)</sup>.

The crude seeds extract of *M. oleifera* was also found to have protective effects against, the toxicity of lead nanoparticles in rats ( Unpublished results), oxidative DNA damage<sup>(19)</sup>, and against bacterial growth<sup>(20)</sup>


## Conclusions

The result of a GC-mass analysis of Iraqi *M. oleifera* was valuable. The number and quality of the compounds manufactured and stored in the seeds of *M. oleifera* reveal the nutritional and/or medicinal importance of these phytochemicals. The compounds, pentadecanoic acid, Cis-Vaccenic acid, dodecanoic acid, and myristicin constitute about 74.64% of the total contents of extraction.




**Figure (1): TIC of the GC-MSS of methanol: acetone (1:1) seeds extract of Iraqi *M.oleifera***

Hit#:2 Entry:37711 Library:NIST08.LIB  
 SI:77 Formula:C<sub>11</sub>H<sub>12</sub>O<sub>3</sub> CAS:607-91-0 MolWeight:192 RetIndex:1516  
 CompName:1,3-Benzodioxole, 4-methoxy-6-(2-propenyl)- \$\$\$\$ Benzene, 5-allyl-1-methoxy-2,3-(methylenedioxy)- \$\$\$\$ Myristicin \$\$\$\$ 5-Allyl-1-methoxy




### A- Myristicin

Hit#:1 Entry:47987 Library:NIST08.LIB  
 SI:82 Formula:C<sub>12</sub>H<sub>16</sub>O<sub>3</sub> CAS:2883-98-9 MolWeight:208 RetIndex:1568  
 CompName:Asarone \$\$\$\$ (E)-2,4,5-Trimethoxypropenylbenzene \$\$\$\$ trans-2,4,5-Trimethoxypropenylbenzene \$\$\$\$ Benzene, 1,2,4-trimethoxy-5-(1-propenyl)




### B- Asarone

Hit#:1 Entry:39106 Library:NIST08.LIB  
 SI:75 Formula:C<sub>11</sub>H<sub>14</sub>O<sub>3</sub> CAS:6627-88-9 MolWeight:194 RetIndex:1581  
 CompName:Phenol, 2,6-dimethoxy-4-(2-propenyl)- \$\$\$\$ Phenol, 4-allyl-2,6-dimethoxy- \$\$\$\$ Methoxyeugenol \$\$\$\$ 2,6-Dimethoxy-4-allylphenol \$\$\$\$ 4-Allyl



### C- Methoxyeugenol

Hit#:3 Entry:70950 Library:NIST08.LIB  
 SI:92 Formula:C15H30O2 CAS:1002-84-2 MolWeight:242 RetIndex:1869  
 CompName: **Pentadecanoic acid** \$\$ Pentadecyclic acid \$\$ n-Pentadecanoic acid \$\$ n-Pentadecyclic acid \$\$



D- Pentadecanoic acid

**Figure 2 : Some important compounds in methanol: acetone (1:1) seeds extract of Iraqi *M.oleifera***

**Ethical Clearance:** The Research Ethical Committee at scientific research by ethical approval of both environmental and health and higher education and scientific research ministries in Iraq

**Conflict of Interest:** The authors declare that they have no conflict of interest.

**Funding:** Self-funding

### References

1. Naik GH, Priyadarsini KI, Satav JG, Banavalikar MM, Sohoni DP, Biyani MK, Mohan H. Comparative antioxidant activity of individual herbal components used in Ayurvedic medicine. *Phytochemistry*. 2003 May 1;63(1):97-104.
2. Saleh MM, Qader SW, Thaker AA. Gastroprotective activity of *Eruca sativa* leaf extract on ethanol-induced gastric mucosal injury in *Rattus norvegicus*. *Jordan Journal of Biological Sciences*. 2016 Mar;147(3384):1-6.
3. Asraa Adnan Shaker, Abid Ali Thaker and AbdalwahabBdewi Hussain. Protective activity of *capparis spinosa* against aflatoxin b1 poisons induced histopathological changes in mice. *Biochem. Cell. Arch*. 2019, Vol. 19, No. 1, pp. 717-723.
4. World Health Organization. Report WHO Health Benefit of Herbs. The Past, the present, the future. *Medicinal Journal of USA*. 2003;182(4):1-0.
5. Padayachee B, Baijnath H. An overview of the medicinal importance of *Moringaceae*. *Journal of Medicinal Plants Research*. 2012 Dec 17;6(48):5831-9.
6. Saini RK. "Studies on enhancement of carotenoids, folic acid, iron and their bioavailability in *Moringa oleifera* and in vitro propagation (Doctoral dissertation, University of Mysore).2013.
7. Compaoré WR, Nikiéma PA, Bassolé HI, Savadogo A, Mouecoucou J. Chemical composition and antioxidative properties of seeds of *Moringa oleifera* and pulps of *Parkia biglobosa* and *Adansonia digitata* commonly used in food fortification in Burkina Faso. *Current Research Journal of Biological Sciences*. 2011 Jan 15;3(1):64-72.
8. Mbah BO, Eme PE, Ogbusu OF. Effect of cooking methods (boiling and roasting) on nutrients and anti-nutrients content of *Moringa oleifera* seeds. *Pakistan Journal of Nutrition*. 2012 Mar 1;11(3):211.
9. Saa RW, Fombang EN, Ndjantou EB, Njintang NY. Treatments and uses of *Moringa oleifera* seeds in human nutrition: A review. *Food science & nutrition*. 2019 Jun;7(6):1911-9.
10. Donli PO, Dauda H. Evaluation of aqueous *Moringa* seed extract as a seed treatment biofungicide for groundnuts. *Pest Management Science: formerly Pesticide Science*. 2003 Sep;59(9):1060-2.
11. Badr G, Elsayy H, Amalki MA, Alfwuaires M, El-Gerbed MS, Abdel-Moneim AM. Protective effects of myristicin against ulcerative colitis induced by acetic acid in male mice. *Food and Agricultural Immunology*. 2020 Jan 1;31(1):435-46.
12. Geng Y, Li C, Liu J, Xing G, Zhou L, Dong M, Li X, Niu Y. Beta-asarone improves cognitive function by suppressing neuronal apoptosis in the beta-amyloid hippocampus injection rats. *Biological and Pharmaceutical Bulletin*. 2010 May 1;33(5):836-43.
13. Tao H, Ding X, Wu J, Liu S, Sun W, Nie M, Pan

- X, Zou X.  $\beta$ -Asarone Increases Chemosensitivity by Inhibiting Tumor Glycolysis in Gastric Cancer. Evidence-Based Complementary and Alternative Medicine. 2020 Apr 30;2020.
14. Bendre RS, Rajput JD, Bagul SD, Karandikar PS. Outlooks on medicinal properties of eugenol and its synthetic derivatives. Nat Prod Chem Res. 2016;4(212):2.
  15. To NB, Nguyen YT, Moon JY, Ediriweera MK, Cho SK. Pentadecanoic Acid, an Odd-Chain Fatty Acid, Suppresses the Stemness of MCF-7/SC Human Breast Cancer Stem-Like Cells through JAK2/STAT3 Signaling. Nutrients. 2020 Jun;12(6):1663.
  16. Fauser JK, Matthews GM, Cummins AG, Howarth GS. Induction of apoptosis by the medium-chain length fatty acid lauric acid in colon cancer cells due to induction of oxidative stress. Chemotherapy. 2013;59(3):214-24.
  17. Abdullah FO, Hussain FH, Mannucci B, Lappano R, Tosi S, Maggiolini M et al. Composition, antifungal and antiproliferative activities of the hydro distilled oils from leaves and flower heads of *Pteroccephalusnestorianus* Nab. Chem Biodivers;2017, 14: e1700009.
  18. AFNS. Alberta natural trans fat research earns global recognition Archived 2008-06-15 at the Wayback Machine. April 2, 2008.
  19. Singh BN, Singh BR, Singh RL, Prakash D, Dhakarey R, Upadhyay G, Singh HB. Oxidative DNA damage protective activity, antioxidant and anti-quorum sensing potentials of *Moringa oleifera*. Food and Chemical Toxicology. 2009 Jun 1;47(6):1109-16.
  20. Emmanuel SA, Olajide OO, Abubakar S, Idowu ID, Orishadipe AT, Thomas SA. Phytochemical and antimicrobial studies of methanol, ethyl acetate, and aqueous extracts of *Moringa oleifera* seeds. American Journal of Ethnomedicine. 2014;1(5):346-54.