

PAPER • OPEN ACCESS

Study of Cellular Immune Response and Some of the Blood Variables in Children with Celiac Disease

To cite this article: Reem K. Ibrahim *et al* 2021 *J. Phys.: Conf. Ser.* **1818** 012011

View the [article online](#) for updates and enhancements.

You may also like

- [Liposome interaction with macrophages and foam cells for atherosclerosis treatment: effects of size, surface charge and lipid composition](#)

Jinkai Tang, Moumita Rakshit, Huei Min Chua *et al.*

- [Surface code—biophysical signals for apoptotic cell clearance](#)

Mona Biermann, Christian Maueröder, Jan M Brauner *et al.*

- [Murine liver damage caused by exposure to nano-titanium dioxide](#)

Jie Hong and Yu-Qing Zhang



The Electrochemical Society
Advancing solid state & electrochemical science & technology

241st ECS Meeting

Vancouver, BC, Canada. May 29 – June 2, 2022

ECS Plenary Lecture featuring
Prof. Jeff Dahn,
Dalhousie University

Register now!

The banner features the ECS logo, a 'Register now!' button with a checkmark, and a photograph of Prof. Jeff Dahn pointing at a whiteboard. The background of the banner shows the Science World geodesic dome in Vancouver, BC, Canada, with a city skyline and water in the foreground.

Study of Cellular Immune Response and Some of the Blood Variables in Children with Celiac Disease

Reem K. Ibrahim¹, QasimKhaif Abdullah², M. J. Muhaidi³, Raghed Khaled Khammas⁴, Najeeb Mohammed Hussein¹

¹University of Anbar, College of Science, Department of Biology, Iraq.

²Ramadi Teaching Hospital for Gynecology and Childhood, senior of Pediatric Department, Ramadi, Iraq.

³University of Fullujahm College of Applied Sciences, , Ramadi, Iraq.

⁴University of Anbar, College of Woman Education, Department of Chemistry.

E-mail: [*drnajeabdulimirartnas@gmail.com](mailto:drnajeabdulimirartnas@gmail.com)

Abstract. All The study presents the effect of the disease celiac in children on cellular immunity. Some of the variables used for this analysis were neutrophil WBC (PMN) rate PMN phagocytosis factor, size granules and movement control factor (MIF) at different ages of 30, 60, 90, and 120 minutes. No significant differences in celiac disease in children on PMNs. Effect celiac disease in children on phagocytosis of *Candida albicans* killed by the heat that there were significant differences between the first and second treatments in children infected compared with control agree with this study.

Keywords: celiac, diseases, cellular, immunity.

1. Introduction

Celiac sickness can be challenging to analyze due to the fact it suddenly influences individuals. There are extra than 200 regarded celiac disorder manifestations that can also manifest in the belly associated framework or exclusive portions of the body. A few humans create a celiac sickness as a youngster, others as a grown-up. The reason at the back of this is as but obscure.[1] A few humans with celiac contamination have no indicators by means of any stretch of the creativeness but at the equal time take a look at positive on the celiac illness blood test.[2] A couple of others can also have a bad blood test; however, have a nice intestinal biopsy.[3] In any case, all men and women with celiac ailment are in threat for lengthy haul entanglements, regardless of whether or not they exhibit any facet effects.[4] Individuals with non-celiac wheat affectability journey manifestations



like these of celiac sickness, which settle when gluten is expelled from the consuming routine. Be that as it may, they do now not take a look at effective for celiac infection.[5]

The celiac illness is a hereditary, immune gadget circumstance in which consuming gluten (a protein determined in wheat, rye, and grain) makes damage the small digestive tract. Patients need to devour gluten robotically for the Symptoms Assessment Tool to be exact. [6] If a affected person has unexplained aspect consequences or has a relative with celiac ailment, entire the Symptoms Assessment Tool to take a look at whether the affected person has an elevated hazard.

2. Methodology

The find out about covered 30 cases, of which 20 had been contaminated with the disease, divided into ten sufferers aged much less than two years, ten infected humans aged greater than two years, and ten non-infected for comparison, the medical analysis of the disease, consisting of diarrhea, weight loss, and anemia, was once taken into consideration.

3. Lab. Test

Some parameters have been used to acquire this study, are 1 WBC and differential count, Concentration of antibodies IgA, IgM, IgG, and Concentration of complement proteins kind C3 and C4/mg/100.[7,8,9,10]

4. Result and discussion

Table 1 suggests the impact of celiac sickness on adolescents contaminated with the disease, the place the effects confirmed no massive variations between contaminated adolescents and non-infected children, and there is no substantial distinction in age in each treatment. This find out about is regular with PMNs have been handled in youngsters youthful than two years of age seventy-nine whilst in teenagers older than two years had been seventy five in both the manipulate was once seventy seven These consequences have been agreed with the lookup of Bartoloni et. al.[11]

Table 1. Effect celiac sickness in young people on PMNs.

Treatment	PMN Percentage
Children <2years	79.0±0.71 ^a
Children > 2 years	75.2±0.42 ^a
(control)	77.2±0.42 ^a

Letters in the column proper exhibit the giant editions ($p < 0.05$).

Effect celiac disorder in kids on phagocytosis of *Candida albicans* had been killed by means of the warmth as proven in Table 1. The outcomes confirmed that there have been no enormous variations in age for youngsters with this sickness except two years and kids contaminated with this ailment over the two years, as nicely as no widespread variations for the time, the place the consequences confirmed that there is no time variations between all the redress agree with this study[12]. The find out about additionally confirmed that there had been full-size variations between the first and 2d remedies in adolescents contaminated in contrast with manipulate agree with the 13].

Table 2. Effect celiac disorder in teens on PMN migration.

Treatment	migration Zone(mm)	Inhibition factor of migration (MIF)
Children <2years	13.3 ±0.10 ^a	0.64
Children > 2 years	18.3±0.04 ^b	1.00
Control	18.3±0.04 ^b	1.00

Letters in the column proper exhibit the giant editions ($p < 0.05$).

Effect celiac sickness in young people on the migration of PMNs is proven in Table (3). The effects confirmed that there have been considerable variations between the therapy in which teens with the disorder and these beneath the age of two years in contrast with youngsters older than two years and the control, the place the element of inhibition of migration to the first therapy 0.64, the 2d therapy and manipulate amounted to 1. 1 respectively agree with this find out about with Mageret al. [14]

- MIF values less than 0.95: inhibition of migration.
- MIF values above 1.05: stimulation of migration.
- MIF values between 0.95 and 1.05: normal migration.

Table 3. Effect celiac ailment in teens Total protein stage g / one hundred ml \pm Standard error (SE).

Treatment	Total protein (G/100 ml)
Children <2years	12.3 \pm 0.10 ^a
Children >2 years	17.3 \pm 0.04 ^b
Control	17.3 \pm 0.04 ^b

Letters in the column proper exhibit the considerable variants ($p < 0.05$).

5. Conclusion

The results of the present study indicate that there is a significant difference between the first treatment and control, as we note that there is a decrease in the total protein concentration in children with the disease and their ages less than two years compared to control, the serum protein is a small part of the body's proteins, It can be studied from proteins in vivo [15, 16]. There were no significant differences between the treatment of children infected with the disease and those older than two years compared to the first treatment [17].

References

- [1] Mayassi T, Ladell K, Gudjonson H, McLaren J E, Shaw D G, Tran M T, Rokicka J J, Lawrence I, Grenier J C, van Unen V, Ciszewski C, Dimaano M, Sayegh H E, Kumar V, Wijmenga C, Green P H R, Gokhale R, Jericho H, Semrad C E, Guandalini S, Dinner A R, Kupfer S S, Reid H H, Barreiro L B, Rossjohn J, Price D A and Jabri B 2019 Chronic Inflammation Permanently Reshapes Tissue-Resident Immunity in Celiac Disease *Cell* 176 967-81.e19
- [2] Vlasova A N, Takanashi S, Miyazaki A, Rajashekara G and Saif L J 2019 How the gut microbiome regulates host immune responses to viral vaccines *Current Opinion in Virology* 37 16-25
- [3] Ponce de Leon C, Angel Lopez-Casado M, Lorite P, Palomeque T and Isabel Torres M 2019 Dysregulation of the PD-1/PD-L1 pathway contributes to the pathogenesis of celiac disease *Cellular & molecular immunology* 16 777-9
- [4] Tsantsaridou A, Papathanasiou V and Dimisianos G 2019 Association of Food Intolerance with Coronary Artery Disease *Journal of Food and Nutrition Research* 7 71-81
- [5] Lania G, Nanayakkara M, Maglio M, Auricchio R, Porpora M, Conte M, De Matteis M A, Rizzo R, Luini A, Discepolo V, Troncone R, Auricchio S and Barone M V 2019 Constitutive alterations in vesicular trafficking increase the sensitivity of cells from celiac disease patients to gliadin *Communications Biology* 2 190

- [6] Maiuri L, Vilella V R, Piacentini M, Raia V and Kroemer G 2019 Defective proteostasis in celiac disease as a new therapeutic target *Cell Death & Disease* 10 114
- [7] Maiuri L, Vilella V R, Raia V and Kroemer G 2019 The gliadin-CFTR connection: new perspectives for the treatment of celiac disease *Italian Journal of Pediatrics* 45 NA
- [8] Vilella V R, Esposito S, Ferrari E, Monzani R, Tosco A, Rossin F, Castaldo A, Silano M, Marseglia G L, Romani L, Barlev N A, Piacentini M, Raia V, Kroemer G and Maiuri L 2019 Autophagy suppresses the pathogenic immune response to dietary antigens in cystic fibrosis *Cell Death & Disease* 10 258
- [9] Mao R, Wu L, Zhu N, Liu X, Liu R and Li Y 2019 Naked Oat (*Avena nuda* L.) Oligopeptides: Immunomodulatory Effects on Innate and Adaptive Immunity in Mice via Cytokine Secretion, Antibody Production, and Th Cells Stimulation *Nutrients* 11 927
- [10] Laurikka P, Lindfors K, Oittinen M, Huhtala H, Salmi T, Lahdeaho M L, Ilus T, Maki M, Kaukinen K and Kurppa K 2019 Dietary Factors and Mucosal Immune Response in Celiac Disease Patients Having Persistent Symptoms Despite a Gluten-free Diet *Journal of clinical gastroenterology* 53 507-13
- [11] Bartoloni E, Bistoni O, Alunno A, Cavagna L, Nalotto L, Baldini C, Priori R, Fischetti C, Fredi M, Quartuccio L, Carubbi F, Montecucco C, Doria A, Mosca M, Valesini G, Franceschini F, De Vita S, Giacomelli R, Mirabelli G, Bini V, Gabrielli A, Catassi C and Gerli R 2019 Celiac Disease Prevalence is Increased in Primary Sjogren's Syndrome and Diffuse Systemic Sclerosis: Lessons from a Large Multi-Center Study *Journal of clinical medicine* 8
- [12] Kulkarni D H and Newberry R D 2019 Intestinal Macromolecular Transport Supporting Adaptive Immunity *Cellular and molecular gastroenterology and hepatology* 7 729-37
- [13] Osorio C E, Wen N, Mejias J H, Liu B, Reinbothe S, von Wettstein D and Rustgi S 2019 Development of wheat genotypes expressing a glutamine-specific endoprotease from barley and a prolyl endopeptidase from *Flavobacterium meningosepticum* or *Pyrococcus furiosus* as a potential remedy to celiac disease *Functional & integrative genomics* 19 123-36
- [14] Mager D R, Liu A, Marcon M, Harms K, Brill H, Mileski H, Dowhaniuk J, Nasser R, Carroll M W, Persad R and Turner J M 2019 Diet patterns in an ethnically diverse pediatric population with celiac disease and chronic gastrointestinal complaints *Clinical nutrition ESPEN* 30 73-80.
- [15] Scarnato L, Gadermaier G, Volta U, De Giorgio R, Caio G, Lanciotti R and Del Duca S 2019 Immunoreactivity of Gluten-Sensitized Sera Toward Wheat, Rice, Corn, and Amaranth Flour Proteins Treated with Microbial Transglutaminase *Frontiers in microbiology* 10 470.
- [16] Ribeiro M, Picascia S, Rhazi L, Gianfrani C, Carrillo J M, Rodriguez-Quijano M, Branlard G and Nunes F M 2019 Effect of in situ gluten-chitosan interlocked self-assembled supramolecular architecture on rheological properties and functionality of reduced celiac-toxicity wheat flour *Food Hydrocolloids* 90 266-75.
- [17] Caio G, Volta U, Ursini F, Manfredini R and De Giorgio R 2019 Small bowel adenocarcinoma as a complication of celiac disease: clinical and diagnostic features *BMC Gastroenterology* 19 45.