

Acta Crystallographica Section E

Structure Reports**Online**


ISSN 1600-5368

N,N'-Bis(diphenylmethyl)benzene-1,4-diamine

Aeed S. Al-Fahdawi, Hussain A. Al-Kafajy, Mohamad J. Al-Jeboori, Simon J. Coles, Claire Wilson and Herman Potgieter

Acta Cryst. (2014). **E70**, o66

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



Acta Crystallographica Section E: Structure Reports Online is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. The journal publishes over 4000 structures per year. The average publication time is less than one month.

Crystallography Journals **Online** is available from journals.iucr.org

***N,N'*-Bis(diphenylmethyl)benzene-1,4-diamine**Aeed S. Al-Fahdawi,^{a*} Hussain A. Al-Kafajy,^a Mohamad J. Al-Jeboori,^b Simon J. Coles,^c Claire Wilson^d and Herman Potgieter^e^aDepartment of Chemistry, College of Science, University of Babylon, Iraq,^bDepartment of Chemical Engineering and Chemical Technology, Imperial College London, London SW7 2AZ, England, ^cUK National Crystallography Service, Chemistry, Faculty of Natural and Environmental Sciences, University ofSouthampton, Southampton, SO17 1BJ, England, ^dDiamond Light Source, Harwell Science and Innovation Campus, Didcot, Oxfordshire OX11 0DE, England, and ^eSchool of Research, Enterprise and Innovation, Manchester Metropolitan University, Chester Street, Manchester M1 5GD, England

Correspondence e-mail: aeedchemistry@yahoo.co.uk


Received 4 November 2013; accepted 10 December 2013

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.077; wR factor = 0.208; data-to-parameter ratio = 17.2.

The complete molecule of the title compound, $\text{C}_{32}\text{H}_{28}\text{N}_2$, is generated by crystallographic inversion symmetry. The dihedral angles between the central aromatic ring and the pendant adjacent rings are 61.37 (16) and 74.20 (14)°. The N—H group does not participate in hydrogen bonds and there are no aromatic π – π stacking interactions in the crystal.

Related literature

The reduction of the Schiff-base was as described in Higuchi *et al.* (2003) and Higuchi *et al.* (2000). For the use of dendrimers in the formation of new types of organic-metallic hybrid materials, see: Kim *et al.* (2005); for drug generation, see: Basavaraj *et al.* (2009). For related structures, see: Ge & Ng (2006); Yang *et al.* (2007); Xia *et al.* (2007). Data were collected and processed according to Coles & Gale (2012).

**Experimental***Crystal data*

$\text{C}_{32}\text{H}_{28}\text{N}_2$
 $M_r = 440.56$
 Monoclinic, $P2_1/n$
 $a = 14.784$ (2) Å
 $b = 5.5853$ (8) Å
 $c = 14.896$ (2) Å
 $\beta = 107.914$ (8)°

$V = 1170.4$ (3) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.07$ mm⁻¹
 $T = 100$ K
 $0.1 \times 0.09 \times 0.02$ mm

Data collection

Rigaku AFC12 (Right) diffractometer
 Absorption correction: multi-scan (*CrystalClear-SM Expert*; Rigaku, 2012)
 $T_{\min} = 0.345$, $T_{\max} = 1.000$

10305 measured reflections
 2664 independent reflections
 1254 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.125$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.077$
 $wR(F^2) = 0.208$
 $S = 0.97$
 2664 reflections

155 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.33$ e Å⁻³
 $\Delta\rho_{\min} = -0.29$ e Å⁻³

Data collection: *CrystalClear-SM Expert* (Rigaku, 2012); cell refinement: *CrystalClear-SM Expert*; data reduction: *CrystalClear-SM Expert*; program(s) used to solve structure: *SHELXD* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *OLEX2*.

The authors would like to thank the 'Iraqi Ministry for Higher Education' for providing six months funding for Mr Aeed S. Al-Fahdawi's PhD scholarship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB7158).

References

- Basavaraj, B. V., Furtado, F., Bharath, S., Deveswaran, R., Sindhu, A. & Madhavan, V. (2009). *J. Pharm. Res.* **2**, 970–974.
 Coles, S. J. & Gale, P. A. (2012). *Chem. Sci.* **3**, 683–689.
 Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
 Ge, W.-Z. & Ng, S. W. (2006). *Acta Cryst.* **E62**, o3784–o3785.
 Higuchi, M., Shiki, S. & Yamamoto, K. (2000). *Org. Lett.* **2**, 3079–3082.
 Higuchi, M., Tsuruta, M., Chiba, H., Shiki, S. & Yamamoto, K. (2003). *J. Am. Chem. Soc.* **125**, 9988–9997.
 Kim, Y.-G., Garcia-Martinez, J. C. & Crooks, R. M. (2005). *Langmuir*, **21**, 5485–5491.
 Rigaku (2012). *CrystalClear-SM Expert*. Rigaku Americas Corporation, The Woodlands, Texas, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Xia, H.-T., Liu, Y.-F., Yang, S.-P. & Wang, D.-Q. (2007). *Acta Cryst.* **E63**, o40–o41.
 Yang, S.-P., Li-Jun, H., Da-Qi, W. & Tie-Zhu, D. (2007). *Acta Cryst.* **E63**, o244–o246.

supplementary materials

Acta Cryst. (2014). E70, o66 [doi:10.1107/S1600536813033497]

***N,N'*-Bis(diphenylmethyl)benzene-1,4-diamine**

Aeed S. Al-Fahdawi, Hussain A. Al-Kafajy, Mohamad J. Al-Jeboori, Simon J. Coles, Claire Wilson and Herman Potgieter

1. Comment

Bis-amine compounds are essential building blocks to produce branched or dendritic polymers. Dendrimers are an interesting class of materials which are based on bis-aromatic imine and amine precursors. These polymeric materials have attracted increasing attention due to their functional coordination groups, which can trap many metal ions or metal clusters within the voids in the dendrimers. This can lead to the formation of new types of organic-metallic hybrid nanomaterials (Kim *et al.*, 2005). Furthermore, the polyvalent nature of dendrimers is a key factor in generating a new class of drugs with much improved and acceptable pharmacokinetic profiles (Basavaraj *et al.*, 2009). This paper reports on a new addition to the bis-amine compounds and its chemical and physical features.

The compound, with a molar mass of 440.56 g mol⁻¹, crystallizes in a monoclinic crystal structure with a space group notation of *P2₁/n* and had a calculated density of 1.250 g cm⁻³. The asymmetric unit consists of half the molecule, the molecule is completed by inversion symmetry. Infrared spectra indicates typical absorbance bands of the functional phenyl group and amine –C=N group at 1570 and 1620 cm⁻¹, respectively. The positive ES mass spectrum of the bis-amine showed a parent ion peak at *m/z* = 441.2362 (*M*+H)⁺, corresponding to C₃₂H₂₈N₂, for which the required value = 440.2252.

2. Experimental

The bis-amine {N1,N4-dibenzhydrylbenzene-1,4-diamine} was prepared in a two-step procedure as follows: (i) A Schiff-base {N1,N4-bis-(diphenylmethylene)benzene-1,4-diamine} was synthesized by adopting a conventional procedure (Higuchi *et al.*, 2000) as follows: A mixture of benzophenone (1.69 g, 9.25 mmol), *p*-phenylenediamine (0.500 g, 4.62 mmol), and 1,4-diaza-bicyclo-[2.2.2]octane (DABCO) (3.11 g, 27.7 mmol) in chlorobenzene (40 ml) was stirred at room temperature for 10 min. Titanium (IV) tetrachloride (1.32 g, 6.93 mmol) dissolved in chlorobenzene (10 ml) was added dropwise using a pressure-equalized dropping funnel. The reaction mixture was heated in an oil bath at 125 °C for 24 h. The precipitate was removed by filtration, and then the filtrate was concentrated. The Schiff-base product (yield: 1.83 g, 91%) was isolated by silica gel uniplat chromatography with an eluent mixture of hexane:ethylacetate; 9:1, R_f = 0.25. (ii) The reduction of the Schiff-base was achieved using conventional procedures (Higuchi *et al.*, 2000; 2003) as follows: NaBH₄ (0.06 g, 1.74 mmol) was added cautiously and in small portions to a mixture of the Schiff-base {N1,N4-bis-(diphenylmethylene)benzene-1,4-diamine} (0.500 g, 0.437 mmol), and SnCl₂ (0.17 g, 0.87 mmol) dissolved in a mixture of dichloromethane/acetonitrile (1:1) (200 ml). The reaction mixture was stirred at room temperature for 10 min under an Argon atmosphere. The crude mixture was washed with an aqueous solution of 1% triethylamine (4x100), and the organic layer was dried over Na₂SO₄. The secondary bis-amine was purified from the crude product by uniplat silica gel chromatography with eluent (hexane: acetonitrile: chloroform; 8: 2: 1), R_f = 0.5. Yield: 0.98 g, 54.14%. Colourless plates were obtained from slow evaporation of a methanol solution of the bis-amine in air.

3. Refinement

Data were collected and processed according to Coles & Gale (2012). Hydrogen atoms were placed in geometrically calculated positions and included as part of a riding model with U_{iso} values set at 1.2 times U_{eq} of the parent atom.

Computing details

Data collection: *CrystalClear-SM Expert* (Rigaku, 2012); cell refinement: *CrystalClear-SM Expert* (Rigaku, 2012); data reduction: *CrystalClear-SM Expert* (Rigaku, 2012); program(s) used to solve structure: *SHELXD* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *OLEX2* (Dolomanov *et al.*, 2009).




Figure 1

The structure of the title compound displacement ellipsoids drawn at the 50% probability level. Symmetry code: (i) $1 - x, 1 - y, -z$.

N,N'-Bis(diphenylmethyl)benzene-1,4-diamine

Crystal data

$\text{C}_{32}\text{H}_{28}\text{N}_2$
 $M_r = 440.56$
 Monoclinic, $P2_1/n$
 $a = 14.784$ (2) Å
 $b = 5.5853$ (8) Å
 $c = 14.896$ (2) Å
 $\beta = 107.914$ (8)°
 $V = 1170.4$ (3) Å³
 $Z = 2$

$F(000) = 468$
 $D_x = 1.250$ Mg m⁻³
 Mo $K\alpha$ radiation, $\lambda = 0.71075$ Å
 Cell parameters from 5636 reflections
 $\theta = 3.4\text{--}27.5^\circ$
 $\mu = 0.07$ mm⁻¹
 $T = 100$ K
 Plate, colourless
 $0.1 \times 0.09 \times 0.02$ mm

Data collection

Rigaku AFC12 (Right)
 diffractometer
 Radiation source: Rotating Anode
 Confocal monochromator
 Detector resolution: 28.5714 pixels mm⁻¹
 profile data from ω -scans

Absorption correction: multi-scan
 (*CrystalClear-SM Expert*; Rigaku, 2012)
 $T_{\text{min}} = 0.345, T_{\text{max}} = 1.000$
 10305 measured reflections
 2664 independent reflections
 1254 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.125$

$$\theta_{\max} = 27.5^\circ, \theta_{\min} = 3.4^\circ$$

$$h = -19 \rightarrow 18$$

$$k = -7 \rightarrow 6$$

$$l = -16 \rightarrow 19$$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.077$
 $wR(F^2) = 0.208$
 $S = 0.97$
 2664 reflections
 155 parameters
 0 restraints
 Primary atom site location: dual
 Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0886P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.33 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.29 \text{ e } \text{Å}^{-3}$
 Extinction correction: *SHELXL97* (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001xFc^2/\lambda^3/\sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.026 (5)

Special details

Experimental. FT—IR data were recorded on a Nicolet ATR FT—IR, while NMR data were collected on a Bruker 400 MHz spectrometer in CD_2Cl_2 -d2 solutions. The assignment of the chemical shifts for the NMR data were made following numbering shown in structure B. Schiff-base {N1,N4-bis(diphenylmethylene)benzene-1,4-diamine}; IR (ATR cm⁻¹) 1620 (C=N), 1597 and 1570 (phenyl). NMR data (p.p.m.), δH (400 MHz, CD_2Cl_2): 6.47 (4H, m; C3, 3', 11, 11'-H), 7.06 (2H, d, J = 7.33 Hz; C15, 15'-H), 7.73 (2H, d, J = 7.33 Hz; C16, 16'-H), 7.27–7.40 (20H, m; aromatic-H); δC (100.63 MHz, CD_2Cl_2): 121.53–136.75, (aromatic carbon); 140.12 (C6, 6', 8, 8'); 147.37 (C14, 14'); 168.24 (C7, 7'); DEPT 13 C NMR exhibited no signals between 140–170 p.p.m.. The positive ES mass spectrum of the bis-amine showed the parent ion peak at $m/z = 441.2362$ ($M+H$)⁺ (95%) corresponding to C₃₂H₂₈N₂; required value = 440.2252. Peaks detected at $m/z = 247.16$ (100%) and 167.09 (98%), correspond to $[M-(\text{ph})_2\text{CH}_2]^+$ and $[M-(\text{ph})_2\text{CH}_2+\text{H}_2\text{N}_2\text{ph}]^+$, respectively.

bis-amine {N1,N4-dibenzhydrylbenzene-1,4-diamine IR (ATR cm⁻¹): 3392 (N—H), 2932; 2873 (C—H) aliphatic, 1599 and 1510 (phenyl). NMR data (p.p.m.), δH (400 MHz, CD_2Cl_2): 3.95 (2H, s, Na, a'-H), 5.36 (2H, s; C7, 7'-H), 6.37 (4H, d, J=7.33 Hz; C15, 15', 16, 16'-H), 7.21–7.36 (20H, m, Ar—H); δC (100.63 MHz, CD_2Cl_2): 49.10 (C7, 7'); 115.21 (C15, 15', 16, 16'); 127.25–129.04 (aromatic carbon); 140 (C6, 6', 8, 8'); 144.07 (C14, 14'), DEPT 13 C NMR exhibited no signals between 140–145 p.p.m.. The positive ES mass spectrum of the bis-amine showed the parent ion peak at $m/z = 441.2362$ ($M+H$)⁺ (95%) corresponding to C₃₂H₂₈N₂; required value = 440.2252. Peaks detected at $m/z = 247.16$ (100%) and 167.09 (98%), correspond to $[M-(\text{ph})_2\text{CH}_2]^+$ and $[M-(\text{ph})_2\text{CH}_2+\text{H}_2\text{N}_2\text{ph}]^+$, respectively.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.44891 (16)	0.4954 (4)	0.1672 (2)	0.0366 (7)
H1	0.4144	0.6110	0.1765	0.044*
C2	0.47809 (19)	0.3103 (5)	0.2375 (2)	0.0294 (8)
H2	0.4635	0.1559	0.2050	0.035*
C3	0.58440 (19)	0.3135 (5)	0.2909 (2)	0.0273 (7)
C4	0.6426 (2)	0.5024 (5)	0.2829 (2)	0.0295 (8)
H4	0.6169	0.6316	0.2440	0.035*
C5	0.7386 (2)	0.4993 (5)	0.3323 (2)	0.0314 (8)

H5	0.7768	0.6270	0.3265	0.038*
C6	0.7785 (2)	0.3094 (5)	0.3901 (2)	0.0326 (8)
H6	0.8431	0.3080	0.4231	0.039*
C7	0.7208 (2)	0.1209 (5)	0.3983 (2)	0.0329 (8)
H7	0.7468	-0.0081	0.4371	0.039*
C8	0.6248 (2)	0.1228 (5)	0.3492 (2)	0.0325 (8)
H8	0.5868	-0.0051	0.3554	0.039*
C9	0.41899 (19)	0.3290 (5)	0.3050 (2)	0.0303 (8)
C10	0.4294 (2)	0.5250 (5)	0.3644 (3)	0.0392 (9)
H10	0.4723	0.6451	0.3626	0.047*
C11	0.3771 (2)	0.5438 (6)	0.4259 (3)	0.0433 (9)
H11	0.3845	0.6767	0.4651	0.052*
C12	0.3137 (2)	0.3662 (6)	0.4296 (3)	0.0416 (9)
H12	0.2782	0.3785	0.4712	0.050*
C13	0.3032 (2)	0.1706 (6)	0.3712 (3)	0.0421 (10)
H13	0.2614	0.0487	0.3742	0.051*
C14	0.3545 (2)	0.1546 (5)	0.3081 (3)	0.0386 (9)
H14	0.3454	0.0245	0.2673	0.046*
C15	0.47494 (18)	0.4941 (5)	0.0834 (2)	0.0283 (8)
C16	0.5290 (2)	0.3109 (5)	0.0627 (2)	0.0315 (8)
H16	0.5489	0.1834	0.1042	0.038*
C17	0.44676 (19)	0.6813 (5)	0.0199 (2)	0.0294 (8)
H17	0.4107	0.8049	0.0332	0.035*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0298 (14)	0.0449 (16)	0.039 (2)	0.0124 (13)	0.0167 (14)	0.0079 (14)
C2	0.0222 (15)	0.0330 (16)	0.032 (2)	0.0024 (13)	0.0070 (14)	-0.0015 (14)
C3	0.0212 (14)	0.0322 (16)	0.030 (2)	-0.0007 (13)	0.0109 (14)	-0.0032 (14)
C4	0.0244 (15)	0.0290 (15)	0.035 (2)	0.0028 (13)	0.0094 (14)	0.0020 (14)
C5	0.0255 (15)	0.0340 (17)	0.035 (2)	-0.0023 (14)	0.0094 (15)	-0.0016 (15)
C6	0.0218 (15)	0.0415 (18)	0.034 (2)	0.0026 (14)	0.0076 (15)	-0.0063 (15)
C7	0.0279 (16)	0.0329 (17)	0.039 (2)	0.0084 (14)	0.0125 (16)	0.0026 (15)
C8	0.0257 (16)	0.0318 (17)	0.042 (2)	0.0006 (13)	0.0137 (16)	0.0046 (15)
C9	0.0210 (14)	0.0349 (17)	0.035 (2)	0.0010 (14)	0.0089 (14)	0.0025 (15)
C10	0.0319 (17)	0.0354 (18)	0.053 (3)	-0.0069 (15)	0.0177 (18)	-0.0066 (17)
C11	0.0378 (19)	0.047 (2)	0.048 (3)	0.0018 (17)	0.0177 (18)	-0.0068 (18)
C12	0.0345 (18)	0.050 (2)	0.048 (3)	0.0093 (17)	0.0248 (18)	0.0082 (18)
C13	0.0324 (18)	0.049 (2)	0.053 (3)	-0.0051 (16)	0.0253 (18)	0.0042 (18)
C14	0.0308 (17)	0.0357 (17)	0.053 (3)	-0.0058 (15)	0.0187 (17)	-0.0035 (16)
C15	0.0156 (13)	0.0345 (16)	0.032 (2)	-0.0022 (13)	0.0039 (14)	-0.0001 (15)
C16	0.0219 (14)	0.0351 (17)	0.037 (2)	0.0020 (13)	0.0086 (14)	0.0049 (15)
C17	0.0182 (14)	0.0350 (17)	0.036 (2)	0.0025 (13)	0.0106 (14)	-0.0022 (15)

Geometric parameters (\AA , $^\circ$)

N1—H1	0.8600	C9—C10	1.386 (4)
N1—C2	1.440 (4)	C9—C14	1.373 (4)
N1—C15	1.415 (4)	C10—H10	0.9300

C2—H2	0.9800	C10—C11	1.372 (4)
C2—C3	1.528 (4)	C11—H11	0.9300
C2—C9	1.525 (4)	C11—C12	1.378 (4)
C3—C4	1.389 (4)	C12—H12	0.9300
C3—C8	1.388 (4)	C12—C13	1.375 (5)
C4—H4	0.9300	C13—H13	0.9300
C4—C5	1.384 (4)	C13—C14	1.380 (4)
C5—H5	0.9300	C14—H14	0.9300
C5—C6	1.379 (4)	C15—C16	1.391 (4)
C6—H6	0.9300	C15—C17	1.385 (4)
C6—C7	1.384 (4)	C16—H16	0.9300
C7—H7	0.9300	C16—C17 ⁱ	1.384 (4)
C7—C8	1.382 (4)	C17—C16 ⁱ	1.384 (4)
C8—H8	0.9300	C17—H17	0.9300
C2—N1—H1	118.8	C10—C9—C2	120.1 (2)
C15—N1—H1	118.8	C14—C9—C2	121.2 (3)
C15—N1—C2	122.4 (2)	C14—C9—C10	118.7 (3)
N1—C2—H2	107.6	C9—C10—H10	119.6
N1—C2—C3	113.6 (2)	C11—C10—C9	120.8 (3)
N1—C2—C9	109.0 (2)	C11—C10—H10	119.6
C3—C2—H2	107.6	C10—C11—H11	120.0
C9—C2—H2	107.6	C10—C11—C12	120.1 (3)
C9—C2—C3	111.2 (3)	C12—C11—H11	120.0
C4—C3—C2	121.9 (3)	C11—C12—H12	120.3
C8—C3—C2	119.5 (2)	C13—C12—C11	119.5 (3)
C8—C3—C4	118.6 (3)	C13—C12—H12	120.3
C3—C4—H4	119.8	C12—C13—H13	119.9
C5—C4—C3	120.4 (3)	C12—C13—C14	120.3 (3)
C5—C4—H4	119.8	C14—C13—H13	119.9
C4—C5—H5	119.6	C9—C14—C13	120.6 (3)
C6—C5—C4	120.9 (3)	C9—C14—H14	119.7
C6—C5—H5	119.6	C13—C14—H14	119.7
C5—C6—H6	120.6	C16—C15—N1	122.1 (3)
C5—C6—C7	118.9 (3)	C17—C15—N1	119.5 (3)
C7—C6—H6	120.6	C17—C15—C16	118.5 (3)
C6—C7—H7	119.7	C15—C16—H16	120.1
C8—C7—C6	120.6 (3)	C17 ⁱ —C16—C15	119.9 (3)
C8—C7—H7	119.7	C17 ⁱ —C16—H16	120.1
C3—C8—H8	119.7	C15—C17—H17	119.2
C7—C8—C3	120.7 (3)	C16 ⁱ —C17—C15	121.7 (3)
C7—C8—H8	119.7	C16 ⁱ —C17—H17	119.2
N1—C2—C3—C4	10.4 (4)	C4—C5—C6—C7	-0.2 (5)
N1—C2—C3—C8	-169.4 (3)	C5—C6—C7—C8	0.1 (4)
N1—C2—C9—C10	-67.2 (4)	C6—C7—C8—C3	0.0 (5)
N1—C2—C9—C14	112.5 (3)	C8—C3—C4—C5	-0.1 (4)
N1—C15—C16—C17 ⁱ	-179.5 (3)	C9—C2—C3—C4	-113.0 (3)
N1—C15—C17—C16 ⁱ	179.5 (3)	C9—C2—C3—C8	67.2 (3)

C2—N1—C15—C16	1.3 (4)	C9—C10—C11—C12	0.4 (5)
C2—N1—C15—C17	-178.1 (3)	C10—C9—C14—C13	-1.9 (5)
C2—C3—C4—C5	-180.0 (3)	C10—C11—C12—C13	0.0 (5)
C2—C3—C8—C7	179.9 (3)	C11—C12—C13—C14	-1.3 (5)
C2—C9—C10—C11	-179.7 (3)	C12—C13—C14—C9	2.3 (5)
C2—C9—C14—C13	178.4 (3)	C14—C9—C10—C11	0.6 (5)
C3—C2—C9—C10	58.8 (4)	C15—N1—C2—C3	69.6 (3)
C3—C2—C9—C14	-121.5 (3)	C15—N1—C2—C9	-165.8 (2)
C3—C4—C5—C6	0.2 (5)	C16—C15—C17—C16 ⁱ	0.1 (5)
C4—C3—C8—C7	0.1 (4)	C17—C15—C16—C17 ⁱ	-0.1 (4)

Symmetry code: (i) $-x+1, -y+1, -z$.