

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/335205694>

Relationship between vitamin D deficiency and serum ferritin level in healthy women

Article in *International Journal of Research in Pharmaceutical Sciences* · July 2019

DOI: 10.26452/ijrps.v10i3.1431

CITATIONS

2

READS

1,280

7 authors, including:



Mohanad Adel Mohammed

Baghdad Medical City

3 PUBLICATIONS 19 CITATIONS

[SEE PROFILE](#)



Mushtaq Talib Abed

Ministry of Health, Iraq. Anbar health directorate. Ramadi teaching hospital for m...

4 PUBLICATIONS 2 CITATIONS

[SEE PROFILE](#)



Muhannad Shweash

College of Pharmacy, University of Anbar

26 PUBLICATIONS 258 CITATIONS

[SEE PROFILE](#)



Hayder Adnan Fawzi

85 PUBLICATIONS 118 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Evaluation of TWIST Score in Predicting Testicular Torsion in Children [View project](#)



Radiology [View project](#)



Relationship between vitamin D deficiency and serum ferritin level in healthy women

Mohanad Adel Mohammed^{*1}, Mushtaq Talib Abed², Muhannad Shweash³, Hayder Adnan Fawzi⁴, Muntaha Qasim Hendi², Ashwaq Talib Duraij², Ghuran Mahdi Salih²

¹Department of Clinical Pharmacy Head, FICMS, College of Pharmacy, Al Farahidi University, Iraq

²Department of Clinical Pharmacy, Ministry of Health, Anbar Health Directorate, Ramadi Teaching Hospital, Hospital for Maternity and Children, Iraq

³Department of Clinical Laboratory Science, College of Pharmacy, University of Anbar, Iraq

⁴Department of clinical pharmacy, Baghdad Teaching hospital, Medical City Complex, Iraq

Article History:

Received on: 07.03.2019

Revised on: 02.06.2019

Accepted on: 06.06.2019

Keywords:

anemia,
healthy women,
vitamin D3 deficiency,
age

ABSTRACT

Vitamin D is an essential vitamin that regulates many biological processes and involved in the activity of many organs like the skin, bone, kidney, etc. ferritin is an essential marker for assessment of anemic status. The current study aimed to assess the relationship between ferritin and vitamin D3 in healthy women at various age groups. A cross-sectional study carried out in Ramadi city and involved 92 healthy women aged from 20-50 years, the participants divided into two groups according to age: group I with age 20 – 35 years and group II with age 36 – 50 years. In the present study mean vitamin D3 in group II 11.8 ± 3.5 ng/dL was significantly lower than group I (35.3 ± 12.2 ng/dL) p -value < 0.001 , a similar finding observed for ferritin (19.6 ± 13.9 vs 66.7 ± 52.1 , p -value < 0.001). There was a direct relationship between ferritin with vitamin D. However, this relationship was only significant in group II (p -value < 0.05), while in group I it was statistically significant. In conclusion, low vitamin D levels associated with low ferritin, indicating that vitamin D deficiency is associated with anemia.



*Corresponding Author

Name: Mohanad Adel Mohammed

Phone: +9647722627943

Email: mohanad.adel@alfarahidiuc.edu.iq

ISSN: 0975-7538

DOI: <https://doi.org/10.26452/ijrps.v10i3.1431>

Production and Hosted by

Pharmascope.org

© 2019 | All rights reserved.

INTRODUCTION

Vitamin D deficiency show an increase trend globally (YETLEY, 2008; MITHAL *et al.*, 2009), several causes of vitamin D deficiency exist including reduce intake, low absorption, reduction in sun exposure, increased hepatic catabolism, or low endoge-

nous synthesis (via the liver or the kidney) (NAIR and MASEEH, 2012).

Ferritin is a large, 24 subunit protein consisting of light and heavy chains (AROSIO and LEVI, 2010). Ferritin represents the intracellular iron storage in the body. Also, it is an acute phase reactant (WANG *et al.*, 2010). Much of the iron stored in ferritin is accessible for metabolic needs (DOWDLE *et al.*, 2014). When ferritin accumulates, it aggregates and is proteolyzed by lysosomal enzymes; it is then converted to an iron-rich, poorly characterized hemosiderin which releases its iron slowly and is detected in cells by the Prussian blue reaction (VAISMAN *et al.*, 2000). There is a highly notable vitamin D3 deficiency in different age groups, especially in females. Vitamin D3 may affect many other parameters and functions in the body. The present study was prearranged to assess the correlation between vitamin D3 deficiency and serum ferritin level in women.

METHODS

Study design

The observational cross-sectional study, the study included 92 healthy adult women, the candidates were classified into two groups according to age: group I with age 20 – 35 years and group II with age 36 – 50 years.

Candidate's selection criteria

After obtaining information and eliciting detailed medical and pathological history, only those who fulfilled the inclusion criteria were included in the study. The candidates were overall healthy adults female aged 20-50 years from Ramadi city. The nominees were selected from some of the students in our college, from outpatient attendants in the central laboratory in Ramadi teaching hospital for maternity and children, in addition to some of the highly confident private laboratories. Most of them have signs of vitamin D3 deficiency like muscle ache, tiredness and hair falling. Written informed consent obtained from all the participants in the study.

Inclusion criteria

- Females aged (20 – 55 years)
- Do not receive vitamin D supplementation

Exclusion criteria

- Diabetic patients
- Renal impairment
- Liver disease

Serum Preparation

Blood samples were collected aseptically by venipuncture into a dry clean and sterile white tube without anticoagulant substances and make it clot. Each tube is recognized with written information for each participant. Blood samples allowed to stand for 20-30 min for clot formation and centrifuged. The supernatant serums were taken and stored in Eppendorf tube at (-20 C to - 80 C) for subsequent analysis or use.

Laboratory tests

Measurement of serum vitamin D3 and ferritin level was done by MINI VIDAS® apparatus, which is a high quality compact automated immunoassay system based on Enzyme-Linked Fluorescent Assay (ELFA) technique. It is greatly appreciated worldwide for its simplicity, flexibility and accuracy.

Data analysis

Analysis of the result was made using mean \pm standard deviation and Pearson's correlation coefficient, and one-sample t-test was used wherever applicable. The results were expressed as percentage and significance, by using Graph Pad Prism® Version 5.0 software.

RESULTS AND DISCUSSION

Vitamin D3 and serum ferritin were significantly lower in group II compared to group I, as illustrated in Table 1 .

There was a direct relationship between ferritin with vitamin D. However, this relationship was only significant in the group II (p-value <0.05), while in group I it was statistically significant, as illustrated in Table 2 and Figure 1 .




Figure 1: Scatterplot of the relationship between ferritin with vitamin D₃

Through the last decade, vitamin D has gained considerable interest in health and biomedical research (HOLICK *et al.*, 2011) . Globally, vitamin D deficiency is widespread and is considered a pandemic (CHEN and HOLICK, 2008) . The MENA regions, including Iraq, are not spared from this micronutrient deficiency, these countries have one of the highest frequency of low vitamin D3 in the globe (MAALOUF *et al.*, 2007; CHAKHTOURA *et al.*, 2018) . Among the most common risk factors for low vitamin D include female gender and their clothing style, multi-parity, sedentary lifestyle, urban living and socio-economic status for adults, and longer than average breastfeeding (BASSIL *et al.*, 2013) .

In the present study mean vitamin D3 in the older women group (group II) 11.8 ± 3.5 ng/dL was significantly lower than younger women (group I) 35.3 ± 12.2 ng/dL (p-value < 0.001), similar finding observed for ferritin (19.6 ± 13.9 vs. 66.7 ± 52.1 , p-value <0.001), additionally there was significant direct correlation between ferritin and vitamin D3 in

Table 1: Assessment of vitamin D₃ and ferritin

Variables	Group I	Group II	p-value
Variables	Group I	Group II	p-value
Number	40	52	-
Vitamin D3 (ng/dL), mean ± SD	35.3 ± 12.2	11.8 ± 3.5	<0.001 [S]
Ferritin (ng/mL), mean ± SD	66.7 ± 52.1	19.6 ± 13.9	<0.001 [S]

SD: standard deviation, S: significance

Table 2: Correlation between ferritin with vitamin D₃

Variables	Ferritin			
	Group II		Group I	
	r	p-value	r	p-value
Vitamin D3	0.464	0.001 [S]	0.290	0.070

r: regression coefficient

group II. Similar findings observed in other studies, (SIM *et al.*, 2010) was in agreement with present study in which low vitamin D (<30 ng/dL) associated with elevated Ferritin compared to normal vitamin D (≥ 30 ng/dL) [364 vs. 189 ng/ml, p-value = 0.01] (SIM *et al.*, 2010; SYED *et al.*, 2017) study, in linear regression models, vitamin D insufficiency was associated with increased hepcidin levels (β [SE] = 0.6 [0.2], P = 0.01) and reduced hemoglobin (β [SE] = -0.9 [0.5], P = 0.046), suggesting that the concentrations of 25(OH)D ≥ 30 ng/mL are associated with lower hepcidin and higher hemoglobin levels (SYED *et al.*, 2017).

Vitamin D appears to be associated with anemia; though the mechanism is unknown. One possibility is that vitamin D modulates the level of systemic cytokine production, thus reducing the inflammatory milieu that leads to anemia of chronic disease. Both in vivo and in vitro studies have demonstrated that vitamin D3 reduces cytokine production (BLAZSEK *et al.*, 1996). Also, vitamin D stimulates erythroid precursors, vitamin D receptors have been recognized in bone marrow tissues (REICHEL *et al.*, 1989; NORMAN, 2006).

CONCLUSION

Low vitamin D levels associated with low ferritin, indicating that vitamin D deficiency is associated with anemia

REFERENCES

AROSIO, P., LEVI, S. 2010. Cytosolic and mitochondrial ferritins in the regulation of cellular iron homeostasis and oxidative damage. *Biochim Biophys Acta*, 1800:783–92.

BASSIL, D., RAHME, M., HOTEIT, M., GEL, F. U. L. E. I. H. A. N., *et al.* 2013. Hypovitaminosis D in the Middle East and North Africa: Prevalence, risk factors and impact on outcomes. *Dermatoendocrinol*, 5:274–98.

BLAZSEK, I., FARABOS, C., QUITTET, P., LABAT, M. L., BRINGUIER, A. F., TRIANA, B. K., MACHOVER, D., REYNES, M., MISSET, J. L. 1996. Bone marrow stromal cell defects and 1 alpha,25-dihydroxyvitamin D3 deficiency underlying human myeloid leukemias. *Cancer Detect Prev*, 20:31–42.

CHAKHTOURA, M., RAHME, M., CHAMOUN, N., FULEIHAN, E. L. H. A. J. J., *et al.* 2018. Vitamin D in the Middle East and North Africa. *Bone Rep*, 8(135-146):10–1016.

CHEN, T. C., HOLICK, M. F. 2008. Vitamin D deficiency: a worldwide problem with health consequences. *The American Journal of Clinical Nutrition*, 87(1080):10–1093.

DOWDLE, W. E., NYFELER, B., NAGEL, J., ELLING, R. A., LIU, S., TRIANTAFELLOW, E., MENON, S., WANG, Z., HONDA, A., PARDEE, G., CANTWELL, J., LUU, C., CORNELLA-TARACIDO, I., HARRINGTON, E., FEKKES, P., LEI, H., FANG, Q., DIGAN, M. E., BURDICK, D., POWERS, A. F., HELLIWELL, S. B., D'AQUIN, S., BASTIEN, J., WANG, H., WIEDERSCHAIN, D., KUERTH, J., BERGMAN, P., SCHWALB, D., THOMAS, J., UGWONALI, S., HARBINSKI, F., TALLARICO, J., WILSON, C. J., MYER, V. E., PORTER, J. A., BUSSIÈRE, D. E., FINAN, P. M., LABOW, M. A., MAO, X., HAMANN, L. G., MANNING, B. D., VALDEZ, R. A., NICHOLSON, T., SCHIRLE, M., KNAPP, M. S., KEANEY, E. P., MURPHY, L. O. 2014. Selective VPS34 inhibitor blocks autophagy and uncovers a role for NCOA4 in ferritin degradation and iron

- homeostasis in vivo. *Nat Cell. Biol*, 16:1069–79.
- HOLICK, M. F., BINKLEY, N. C., BISCHOFF-FERRARI, H. A., GORDON, C. M., HANLEY, D. A., HEANEY, R. P., MURAD, M. H., WEAVER, C. M. 2011. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*, 96:1911–30.
- MAALOUF, G., GANNAGE-YARED, M. H., EZZEDINE, J., LARIJANI, B., BADAWI, S., RACHED, A., ZAKROUI, L., MASRI, B., AZAR, E., SABA, E., NAMMARI, R., ADIB, G., SAMRA, A. B. O. U., H., A. L. R. A. W. I., Z., S. A. L. M. A. N., S., E. L. M. U. N. T. A. S. S. E. R., K., T. A. R. S. E. E. N., R., E. L. K. H. A. R. O. U. S. I., W., A. L. L. A. M. K. I., M., A. L. O. T. H. M. A. N., N., A., ALMARZOOK, N., DESSOUKI, E. L., M., S. U. L. A. I. M. A. N. I., R., S. A. L. E. H., J., S. U. H. A. I. L. I., R., A., KHAN, A., DELMAS, P., SEEMAN, E. 2007. Middle East and North Africa consensus on osteoporosis. *J Musculoskelet Neuronal Interact*, 7:131–43.
- MITHAL, A., WAHL, D. A., BONJOUR, J. P., BURCKHARDT, P., DAWSON-HUGHES, B., EISMAN, J. A., FULEIHAN, E. L. H. A. J. J., G., J. O. S. S. E., G., R., LIPS, P., MORALES-TORRES, J. 2009. Global vitamin D status and determinants of hypovitaminosis. *D. Osteoporos Int*, 20:1807–20.
- NAIR, R., MASEEH, A. 2012. Vitamin D: The "sunshine" vitamin. *J Pharmacol Pharmacother*, 3:118–26.
- NORMAN, A. W. 2006. Minireview: vitamin D receptor: new assignments for an already busy receptor. *Endocrinology*, 147:5542–8.
- REICHEL, H., KOEFFLER, H. P., NORMAN, A. W. 1989. The role of the vitamin D endocrine system in health and disease. *N Engl J Med*, 320:980–91.
- SIM, J. J., LAC, P. T., LIU, I. L. A., MEGUERDITCHIAN, S. O., KUMAR, V. A., KUJUBU, D. A., RASGON, S. A. 2010. Vitamin D deficiency and anemia: a cross-sectional study. *Annals of Hematology*, 89:447–452.
- SYED, S., MICHALSKI, E. S., TANGPRICHA, V., CHESDACHAI, S., KUMAR, A., PRINCE, J., ZIEGLER, T. R., SUCHDEV, P. S., KUGATHASAN, S. 2017. Vitamin D Status Is Associated with Hepcidin and Hemoglobin Concentrations in Children with Inflammatory Bowel Disease. *Inflamm Bowel Dis*, 23(1650-1658):10–1097.
- VAISMAN, B., MEYRON-HOLTZ, E. G., FIBACH, E., KRICHEVSKY, A. M., KONIJN, A. M. 2000. Ferritin expression in maturing normal human erythroid precursors. *Br J Haematol*, 110:394–401.
- WANG, W., KNOVICH, M. A., COFFMAN, L. G., TORTI, F. M., TORTI, S. V. 2010. Serum ferritin: Past, present and future. *Biochim Biophys Acta*, 1800:760–9.
- YETLEY, E. A. 2008. Assessing the vitamin D status of the US population. *Am J Clin Nutr*, 88(558):10–1093.